Engineering Mathematics IV

One Shot

STATISTICAL TECHNIQUES-II



UNIT-4

WATCH NOW

ONE SHOT PDF NOTES

- ✓ Available in Gateway Classes App
- ✓ Download Gateway Classes App from the Google play store
- ✓ App link in Description
- ✓ DATA STRUCTURE / COA / PYTHON / CYBER SECURITY / UHVPE / TC Playlist link in Description

New channel

Gateway Classes official AKTU

ENGINEERING MATHEMATICS

UNIT-4 : *Statistical Techniques – II*

Lec-01

Today's Target

- ***Random variables(PART-1)***
- PYQ
- DPP

Gateway Classes : 7455 961284

UNIT – 4 : Statistical Technique – II (Probability)

Aktu Syllabus

- **Random variables**

(i) **Discrete random variables**

- **Probability Mass Function**
- **Expectation and Variance**

(ii) **Continous random variables**

- **Probability Density Function**
- **Expectation and Variance**

- **Probability Distribution**

(i) **Binomial** → **Based on Discrete random variable**

(ii) **Poisson** → **Based on Discrete random variable**

(iii) **Normal** → **Based on Continous random variable**

Gateway Classes : 7455 9612 84

Probability

Probability is a concept which numerically meansures the degree of uncertainity and certanity of the occurrence of events.

$0 \leq P \leq 1$

$$\text{Probability} = \frac{\text{Favourable Cases}}{\text{Total number of mutually exclusive and equally likely cases}}$$

Random variable

It is a real valued function whose domain is the sample space of a random experiment

It is denoted by X

$X = \text{Number of things}$

$X : S \longrightarrow R$

Domain codomain

$X(s) = x$

Where x is any real number

Gateway Classes : 7453961284

Types of Random variable

(1) Discrete random variable

(2) Continous random variable

Discrete random variable

A random variable which can assume only a finite number of values is called a discrete random variable.

Example: –The number of tails in 3 tosses of a coin is a discrete random variable (X)

$$S = \{HHH, HHT, HTH, THH, TTT, TTH, THT, HTT\}$$

X = Number of Tails

$X(HHH) = 0$

Gateway Classes: 7455 9612 84

$X(HHT, HTH, THH) = 1$

$X(TTH, THT, HTT) = 2$

$X(TTT) = 3$

Hence

$X = 0, 1, 2, 3$

Probabilities

$$P(X=0) = \frac{1}{8}$$

$$P(X=1) = \frac{3}{8}$$

$$P(X=2) = \frac{3}{8}$$

$$P(X=3) = \frac{1}{8}$$

$$P(X=0) + P(X=1) + P(X=2) + P(X=3) =$$

$$\frac{1}{8} + \frac{3}{8} + \frac{3}{8} + \frac{1}{8}$$

$$= \frac{1+3+3+1}{8}$$

$$= \frac{8}{8} = 1$$

Probability Distribution

X	0	1	2	3
P(X)	$\frac{1}{8}$	$\frac{3}{8}$	$\frac{3}{8}$	$\frac{1}{8}$

Gateway Classes: 7455 9612 84

Prabability Mass Function

Let $x_1, x_2, x_3, \ldots\ldots$ be the values of a discrete random variable X and let $p_1, p_2, p_3, \ldots\ldots$ be the corresponding probabilities.

A function $p(x)$ defined by

$$p(X = x_i) = p(x) = \begin{cases} p(x_i) \text{ or } p_i & i = 1, 2, 3, 4 \ldots\ldots \\ 0, & \text{otherwise} \end{cases}$$

is called the probability mass function of the discrete random variable X where

(i) $p(x_i) \geq 0$

(ii) $\sum p(x_i) = 1$

V. Imp

Gateway Classes : 7455 9612 84

Probability distribution of a discrete random variable

X	x_1	x_2	x_3	$\ldots\ldots\ldots$	x_n
$p(x)$	p_1	p_2	p_3	$\ldots\ldots\ldots$	p_n

Gateway Classes : 7455 9612 84

2/3/4....Balls (withdrawn)

- **Successively**
  - **With replacement (Independent events)**
  - **Without Replacement (Dependent events)**
- **Simultaneously** (Dependent)

$$^{n}C_{r} = \frac{n!}{r!\,(n-r)!}$$

Q. 1 ***A coin is tossed three times. if X is a random variable giving the number of ~~tails~~ Heads that appear, make a table showing the probability distribution of X.***

$S = \{HHH, HHT, HTH, THH, TTT, TTH, THT, HTT\}$

X = Number of Heads

$X(TTT) = 0$

$X(TTH, THT, HTT) = 1$

$X(HHT, HTH, THH) = 2$

$X(HHH) = 3$

Hence

$X = 0, 1, 2, 3$

Probabilities corresponding to random variables

$P(X=0) = \frac{1}{8}$

Gateway Classes :7455 961284

$$P(X=1) = \frac{3}{8}$$

$$P(X=2) = \frac{3}{8}$$

$$P(X=3) = \frac{1}{8}$$

X	0	1	2	3
P(X)	$\frac{1}{8}$	$\frac{3}{8}$	$\frac{3}{8}$	$\frac{1}{8}$

Gateway Classes : 7455 961284

Q. 2 *A random variable X has the following probability distribution:*

Value of X, x:	0	1	2	3	4	5	6	7
$p(x)$:	0	k	$2k$	$2k$	$3k$	k^2	$2k^2$	$7k^2+k$

Find (i) k (ii) $P(X<6)$ (iii) $P(X\geq 6)$ (iv) $P(3<X\leq 6)$

(v) Minimum value of n for which $P(X\leq n)>\frac{1}{2}$

Solution

(i) We know that

$$\sum P(x_i)=1$$

$$0+k+2k+2k+3k+k^2+2k^2+7k^2+k=1$$

$$9k+10k^2=1$$

Gateway Classes : 7455 961284

$$10K^2 + 9K - 1 = 0$$

$$10K^2 + 10K - K - 1 = 0$$

$$10K(K+1) - (K+1) = 0$$

$$(K+1)(10K-1) = 0$$

$K = -1$ (Neglect)

$$\boxed{K = \frac{1}{10}}$$

$$P(X<6) = P(X=0) + P(X=1) + P(X=2) + P(X=3) + P(X=4) + P(X=5)$$

$$= 0 + K + 2K + 2K + 3K + K^2$$

$$= K^2 + 8K$$

$$= \left(\frac{1}{10}\right)^2 + 8 \times \frac{1}{10} = \frac{1}{100} + \frac{8}{10}$$

$$\boxed{P(X<6) = \frac{81}{100}}$$

(iii) $P(X \geq 6) = P(X=6) + P(X=7)$

$$= 2K^2 + 7K^2 + K$$

Gateway Classes : 7455961284

$$P(X \geq 6) = 9k^2 + k$$

$$= \frac{9}{100} + \frac{1}{10}$$

$$= \frac{9 + 10}{100}$$

$$\boxed{P(X \geq 6) = \frac{19}{100}}$$

(iv) $P(3 < X \leq 6) =$

$$= P(X=4) + P(X=5) + P(X=6)$$

$$= 3k + k^2 + 2k^2$$

$$P(3 < X \leq 6) = 3k^2 + 3k = \frac{3}{100} + \frac{3}{10}$$

$$\boxed{P(3 < X \leq 6) = \frac{33}{100}}$$

(v) $P(X \leq 1) = 0 + k$

$$P(X \leq 1) = k$$

$$P(X \leq 1) = \frac{1}{10} < \frac{1}{2}$$

$$P(X \leq 3) = 0 + k + 2k + 2k$$

$$P(X \leq 3) = 5k$$

$$P(X \leq 3) = \frac{5}{10} = \frac{1}{2}$$

$$P(X \leq 4) = 0 + k + 2k + 2k + 3k$$

$$P(X \leq 4) = 8k$$

$$P(X \leq 4) = \frac{8}{10} > \frac{1}{2}$$

Hence Minimum value of n is 4

Gateway Classes : 7455961284

Q.3 ***Five defective bulbs are accidentally mixed with twenty good ones. It is not possible to just look at a bulb and tell wheter or not it is defective. Find the probability distribution of the number of defective bulbs, if four bulbs are drawn at random from this lot.***

Given

Number of Defective bulbs = 5

Number of good bulbs = 20

Total number of bulbs = 25

X = Number of Defective bulbs

X = 0, 1, 2, 3, 4

Probabilities

$$P(X=0) = \frac{{}^{20}C_4}{{}^{25}C_4}$$

5 D
20 G

$$P(X=0) = \frac{20!}{4! \times 16!} \times \frac{4! \times 21!}{25!}$$

$$= \frac{20 \times 19 \times 18 \times 17 \times 16! \times 21!}{16! \times 25 \times 24 \times 23 \times 22 \times 21!}$$

5 4 2 11

Gateway Classes : 7455961284

$$P(X=0) = \frac{19 \times 3 \times 17}{5 \times 2 \times 11 \times 23}$$

$$P(X=0) = \frac{969}{2530}$$

$$P(X=1) = \frac{{}^{5}C_{1} \times {}^{20}C_{3}}{{}^{25}C_{4}}$$

$$P(X=1) = \frac{1140}{2530}$$

$$P(X=2) = \frac{{}^{5}C_{2} \times {}^{20}C_{2}}{{}^{25}C_{4}}$$

$$P(X=2) = \frac{380}{2530}$$

$$P(X=3) = \frac{{}^{5}C_{3} \times {}^{20}C_{1}}{{}^{25}C_{4}}$$

$$P(X=3) = \frac{40}{2530}$$

$$P(X=4) = \frac{{}^{5}C_{4}}{{}^{25}C_{4}}$$

$$P(X=4) = \frac{1}{2530}$$

X	0	1	2	3	4
P(X)	$\frac{969}{2530}$	$\frac{1140}{2530}$	$\frac{380}{2530}$	$\frac{40}{2530}$	$\frac{1}{2530}$

Gateway Classes : 7455 96 12 84

Q.4 *A random variable X takes values 1, 2, 3, ... with probability mass function* $\frac{\lambda^r}{r!}$, $r = 1, 2, 3 \ldots \infty$. *Find the value of* λ.

Given

$$P(x_i) = \frac{\lambda^x}{x!}$$

$$\sum P(x_i) = 1$$

$$\sum \frac{\lambda^x}{x!} = 1$$

$$\frac{\lambda}{1!} + \frac{\lambda^2}{2!} + \frac{\lambda^3}{3!} + \cdots = 1$$

$$1 + \frac{\lambda}{1!} + \frac{\lambda^2}{2!} + \frac{\lambda^3}{3!} + \cdots = 1 + 1$$

$$e^{\lambda} = 2$$

Taking log both side

$$\lambda \log_e e = \log_e 2$$

$$\boxed{\lambda = \log_e 2}$$

Gateway Classes : 7455 9612 84

Topic : *Random variables (PART-1)*

Q.1 *A box has 5 Blue and 3 Red balls. If 2 balls are to be drawn at random without replacement and X denotes the number of Blue balls, find the probability distribution for X.*

Q.2 *A random variable X has the following probability distribution:*

Value of X, x:	0	1	2	3	4	5	6	7	8
$p(x)$:	a	$3a$	$5a$	$7a$	$9a$	$11a$	$13a$	$15a$	$17a$

(*i*) *Determine the value of* a.

(*ii*) *Find* $P(X < 3), P(X \geq 3), P(2 \leq X < 5)$

(*iii*) *What is the smallest value of* x *for which* $P(X \leq x) > 0.5$?

Gateway Classes : 7455 9612 84

ENGINEERING MATHEMATICS

UNIT-4 : *Statistical Techniques – II*

Lec-02

Today's Target

- ***Random variables (PART-2)***
- PYQ
- DPP

Gateway Classes : 7455 961284

Types of Random variable

(1) Discrete random variable

(2) Continous random variable

Discrete random variable

A random variable which can assume only a finite number of values is called a discrete random variable.

- Probability Mass Function
- Probability Distribution
- Expectation and Variance

Gateway Classes : 7455 9612 84

Expectation and Variance

Mean

If X is a discrete random variable with probability distribution

X:	x_1	x_2	x_3	.....	x_n
$p(X)$:	$p(x_1)$	$p(x_2)$	$p(x_3)$	.....	$p(x_n)$

Expectation

$$E(x) = x_1 p(x_1) + x_2 p(x_2) + x_3 p(x_3) + \ldots\ldots\ldots\ldots + x_n p(x_n)$$

$$\boxed{E(x) = \sum x\, p(x)}$$

$$E(x) = \bar{x} = \mu = \mu_1' = \sum x\, p(x)$$

Expectation or Mathematical *Expectation* or *Mean* or *First moment about origin*

Gateway Classes : 7455 9512 84

Important points:

$$E(x) = \sum x\, p(x) = \mu_1' \qquad (First\ moment\ about\ origin)$$

$$E\left(x^2\right) = \sum (x^2) p(x) = \mu_2' \qquad (Second\ moment\ about\ origin)$$

$$E\left(x^3\right) = \sum (x^3) p(x) = \mu_3' \qquad (Third\ moment\ about\ origin)$$

$$E\left(x^r\right) = \sum (x^r) p(x) = \mu_r' \qquad (r^{th} moment\ about\ origin)$$

Properties of Expectation:

$(i)\ E(c) = c \qquad E(1) = 1$

$(ii)\ E(ax) = a\,E(x)$

$(iii)\ E(x \pm y) = E(x) \pm E(y)$

$(iv)\ E(xy) = E(x).E(y)$

Gateway Classes : 7455 961284

Variance

r^{th} moment about mean

$$\mu_r = \frac{\sum p(x-\bar{x})^r}{\sum pi}$$

But $\sum p_i = 1$

$$\mu_\gamma = \sum p(x-\bar{x})^\gamma$$

$$\mu_1 = \sum p(x-\bar{x})$$

$$\mu_1 = E(x-\bar{x})$$

$$\mu_1 = E(x) - E(\bar{x})$$

$$\mu_1 = \bar{x} - \bar{x}$$

$$\mu_1 = 0$$

$$\mu_2 = \sum p(x-\bar{x})^2$$

$$\mu_2 = \sum p(x-\mu)^2$$

$$\text{Variance} = \sum p(x-\mu)^2$$

$$\mu_2 = \sum p(x-\bar{x})^2$$

$$\mu_2 = E(x-\bar{x})^2$$

$$\mu_2 = E(x^2 + \bar{x}^2 - 2x\bar{x})$$

$$\mu_2 = E(x^2) + E(\bar{x}^2) - E(2x\bar{x})$$

$$\mu_2 = E(x^2) + \bar{x}^2 - 2\bar{x}E(x)$$

$$\mu_2 = E(x^2) + \bar{x}^2 - 2\bar{x}\,\bar{x}$$

$$\mu_2 = E(x^2) + \bar{x}^2 - 2\bar{x}^2$$

$$\mu_2 = E(x^2) - (\bar{x})^2$$

$$\mu_2 = E(x^2) - [E(x)]^2$$

μ_2 : Variance OR Second moment about mean

Gateway Classes : 7455 9612 84

Impotant Formulae

1. *Expectation* (*Mean or First moment about origin*)

$$E(x) = x_1 p(x_1) + x_2 p(x_2) + x_3 p(x_3) + \ldots\ldots\ldots\ldots\ldots + x_n p(x_n)$$

$$E(x) = \sum x\, p(x)$$

2. *Variance* (*Second moment about mean*)

$$Variance = \mu_2 = \sum p\,(x-\mu)^2$$

$$Variance = \mu_2 = E(x^2) - [E(x)]^2$$

3. *Standard Deviation*

$$S.D = \sqrt{Variance}$$

$$E(x^2) = x_1^2 P(x_1) + x_2^2 P(x_2) + \ldots + x_n^2 P(x_n)$$

Gateway Classes : 7455 96 12 84

Q.1 Find the Mean, Variance and Standard deviation of the number of heads when three coins are tossed.

$S = \{HHH, HHT, HTH, THH, TTT, TTH, THT, HTT\}$

X = Number of heads

$X(TTT) = 0$

$X(TTH, THT, THH) = 1$

$X(HHT, HTH, THH) = 2$

$X(HHH) = 3$

$X = 0, 1, 2, 3$

Probabilities

$P(X=0) = \frac{1}{8}$

$P(X=1) = \frac{3}{8}$

$P(X=2) = \frac{3}{8}$

$P(X=3) = \frac{1}{8}$

Gateway Classes: 7455 96 12 84

Probability Distribution

X	0	1	2	3
P(X)	$\frac{1}{8}$	$\frac{3}{8}$	$\frac{3}{8}$	$\frac{1}{8}$

Mean

$$E(X) = \bar{x} = 0 \times \frac{1}{8} + 1 \times \frac{3}{8} + 2 \times \frac{3}{8} + 3 \times \frac{1}{8}$$

$$\bar{x} = 0 + \frac{3}{8} + \frac{6}{8} + \frac{3}{8}$$

$$\bar{x} = \frac{12}{8}$$

$$\boxed{\bar{x} = \frac{3}{2}}$$

Variance

$$E(x^2) = 0^2 \times \frac{1}{8} + 1^2 \times \frac{3}{8} + 2^2 \times \frac{3}{8} + 3^2 \times \frac{1}{8}$$

$$E(x^2) = 0 + \frac{3}{8} + \frac{12}{8} + \frac{9}{8}$$

$$E(x^2) = \frac{3+12+9}{8} = \frac{24}{8} = 3$$

$$\text{Variance} = E(x^2) - [E(x)]^2$$

$$= 3 - \left(\frac{3}{2}\right)^2 = 3 - \frac{9}{4}$$

$$\boxed{\text{Variance} = \frac{3}{4}}$$

$$\boxed{SD = \frac{\sqrt{3}}{2}}$$

Gateway Classes: 7455 961284

Q.2 Two cards are drawn simultaneously from a well shuffled pack of 52 cards. Find the Expectation and Variance of the number of kings.

Let X = Number of kings

$X = 0, 1, 2$

$$P(X=0) = \frac{{}^{48}C_2}{{}^{52}C_2}$$

$$= \frac{48!}{2! \times 46!} \times \frac{2! \times 50!}{52!}$$

$$= \frac{48 \times 47 \times 46! \times 50!}{46! \times 52 \times 51 \times 50!}$$

$$P(X=0) = \frac{188}{221}$$

$$P(X=1) = \frac{{}^{4}C_1 \times {}^{48}C_1}{{}^{52}C_2}$$

$$P(X=1) = \frac{32}{221}$$

$$P(X=2) = \frac{{}^{4}C_2}{{}^{52}C_2}$$

$$P(X=2) = \frac{1}{221}$$

$${}^{n}C_r = \frac{n!}{r!\,(n-r)!}$$

Gateway Classes: 7455961284

Probality Distribution

X	0	1	2
P(X)	$\frac{188}{221}$	$\frac{32}{221}$	$\frac{1}{221}$

Expectation

$$E(X) = 0 \times \frac{188}{221} + 1 \times \frac{32}{221} + \frac{2}{221}$$

$$\boxed{E(X) = \frac{2}{13}}$$

$$E(X^2) = 0^2 \times \frac{188}{221} + 1^2 \times \frac{32}{221} + 2^2 \times \frac{1}{221}$$

$$E(X^2) = \frac{36}{221}$$

Variance

$$Var(X) = E(X^2) - [E(X)]^2$$

$$= \frac{36}{221} - \left(\frac{2}{13}\right)^2$$

$$\boxed{Var(X) = 0.1392}$$

Standard deviation

$$S.D = \sqrt{0.1392}$$

$$\boxed{S.D = 0.37}$$

Gateway Classes : 7455 9612 84

Q.3 Let X denote the sum of the numbers obtained when two fair dice are rolled. Find the Expectation of X

X = sum of numbers on two dice

X	2	3	4	5	6	7	8	9	10	11	12
P(X)	$\frac{1}{36}$	$\frac{2}{36}$	$\frac{3}{36}$	$\frac{4}{36}$	$\frac{5}{36}$	$\frac{6}{36}$	$\frac{5}{36}$	$\frac{4}{36}$	$\frac{3}{36}$	$\frac{2}{36}$	$\frac{1}{36}$

Expectation

$$E(X) = \frac{2}{36} + \frac{6}{36} + \frac{12}{36} + \frac{20}{36} + \frac{30}{36} + \frac{42}{36} + \frac{40}{36} + \frac{36}{36} + \frac{30}{36} + \frac{22}{36} + \frac{12}{36}$$

$$E(X) = 7$$

Gateway Classes : 7455961284

Q.4 A random variable X has the following distribution

X:	-2	-1	0	1	2	3
$p(X)$:	0.1	k	0.2	$2k$	0.3	k

Determine (i) k (ii) Mean (iii) Variance

(i) We know that

$$\sum p(x) = 1$$

$$0.1 + k + 0.2 + 2k + 0.3 + k = 1$$

$$4k + 0.6 = 1$$

$$4k = 0.4$$

$$\boxed{k = 0.1}$$

Mean

$$\bar{x} = -0.2 - k + 0 + 2k + 0.6 + 3k$$

$$\bar{x} = -0.2 - 0.1 + 0.2 + 0.6 + 0.3$$

$$\boxed{\bar{x} = 0.8}$$

Variance

$$E(x^2) = (-2)^2 \times 0.1 + 1^2 \times k + 0 + 1^2 \times 2k + 2^2 \times 0.3 + 3^2 \times k$$

Gateway Classes : 7455 9612 84

$$E(x^2) = 0.4 + k + 2k + 1.2 + 9k$$

$$= 0.4 + 0.1 + 0.2 + 1.2 + 0.9$$

$$E(x^2) = 2.8$$

$$\text{Variance} = E(x^2) - [E(x)]^2$$

$$\text{Variance} = 2.8 - (0.8)^2$$

$$= 2.8 - 0.64$$

$$\boxed{\text{Var}(X) = 2.16}$$

Gateway Classes : 7455 9612 84

Engineering Mathematics by Gulshan Sir

DPP- 02

Topic : *Random variables (PART-2)*

$Q.1$ *Let X denote the number of heads when a pair of coin is tossed. What is the expected value?*

$Q.2$ *Two dice are thrown simultaneously. if X denote the number of sixes, find the expectation of X*

Gateway Classes : 7455 9612 84

ENGINEERING MATHEMATICS

UNIT-4 : *Statistical Techniques – II*

Lec-03

Today's Target

- ***Random variables (PART-3)***
- PYQ
- DPP

Gateway Classes : 7455 961284

Types of Random variable

(1) Discrete random variable

(2) Continous random variable

Discrete random variable


A random variable which can assume only a finite number of values is called a discrete random variable.

- Probability Mass Function
- Probability Distribution
- Expectation and Variance

Gateway Classes : 7455 9612 84

Continous random variable

- A random variable which can assume any real value within an interval



- The interval may be finite or infinite

Probability Density Function

Let X be a continous random variable, then in the interval $[a, b]$, the PDF is defined by

$$P(a \leq X \leq b) = \int_a^b f(x)\,dx$$

Such that

(i) $f(x) \geq 0, \quad \forall\, x$

(ii) $\int_{-\infty}^{\infty} f(x)\,dx = 1$

Gateway Classes : 7455 9612 84

Note: $P((a \le X \le b) = f(x) \text{ or } p(x) = \begin{cases} 0 & x < a \\ \emptyset(x) & a \le x \le b \\ 0 & x > b \end{cases}$

$$e^{-\infty} = \frac{1}{e^{\infty}} = \frac{1}{\infty} = 0$$

Q.1 *If the function $f(x)$ is defined by $f(x) = ce^{-x}, 0 \le x \le \infty$, find the value of c which changes $f(x)$ to a probability density function.*

$f(x) = ce^{-x}, \quad 0 \le x \le \infty$

$f(x)$ is a PDF if

(i) $f(x) \ge 0, \forall x$

(ii) $\int_{-\infty}^{\infty} f(x)\,dx = 1$

$\int_{-\infty}^{\infty} f(x)\,dx = 1$

$\int_{-\infty}^{0} f(x)\,dx + \int_{0}^{\infty} f(x)\,dx = 1$

$0 + \int_{0}^{\infty} ce^{-x}\,dx = 1$

$\left(-ce^{-x}\right)_{0}^{\infty} = 1$

$-c\left[e^{-\infty} - e^{0}\right] = 1$

$-c[0 - 1] = 1$

$\boxed{c = 1}$

$f(x) = ce^{-x}$

$e^{-x} \ge 0$

$c \ge 0$

$\Rightarrow f(x) \ge 0$

Gateway Classes 7455 9612 24

Q.2 *If $f(x)$ has probability density $cx^2, 0 < x < 1$, determine c and find the probability that $\frac{1}{3} < x < \frac{1}{2}$ i.e., $P\left(\frac{1}{3} < x < \frac{1}{2}\right)$.*

$f(x) = cx^2, \quad 0 < x < 1$

Since $f(x)$ is a PDF

$\Rightarrow$ (i) $f(x) \geq 0$

(ii) $\int_{-\infty}^{\infty} f(x)\,dx = 1$

From (ii)

$$\int_{-\infty}^{\infty} f(x)\,dx = 1$$

$$\int_{-\infty}^{0} f(x)\,dx + \int_{0}^{1} f(x)\,dx + \int_{1}^{\infty} f(x)\,dx = 1$$

$$0 + \int_{0}^{1} cx^2\,dx + 0 = 1$$

$$c\left(\frac{x^3}{3}\right)_0^1 = 1$$

$$c\left(\frac{1}{3} - 0\right) = 1$$

$$\frac{c}{3} = 1$$

$$\boxed{c = 3}$$

$$f(x) = 3x^2$$

Gateway Classes : 7455 961 284

$$P\left(\frac{1}{3} < x < \frac{1}{2}\right) = \int_{1/3}^{1/2} f(x)\,dx$$

$$= \int_{1/3}^{1/2} 3x^2\,dx$$

$$= 3\left(\frac{x^3}{3}\right)_{1/3}^{1/2}$$

$$= \frac{1}{8} - \frac{1}{27}$$

$$= \frac{27 - 8}{216}$$

$$\boxed{P\left(\frac{1}{3} < x < \frac{1}{2}\right) = \frac{19}{216}}$$

Gateway Classes : 7455 9612 84

Expectation and variance

If x is a continous random variable, then the expectation of x is

$$E(x) = \int_{-\infty}^{\infty} x\, f(x)\, dx$$

Note : $E(x) = \bar{x} = \mu = \mu_1' = \int_{-\infty}^{\infty} x\, f(x)\, dx$

Expectation or mean or first moment about origin

Also

- $E(x) = \int_{-\infty}^{\infty} x\, f(x)\, dx$ **(First moment about origin)**
- $E(x^2) = \int_{-\infty}^{\infty} x^2\, f(x)\, dx$ **(Second moment about origin)**
- $E(x^r) = \int_{-\infty}^{\infty} x^r\, f(x)\, dx$ **(r^{th} moment about origin)**

Gateway Classes 7455 9612 84

Variance

$$\text{Var}(x) = E(x^2) - [E(x)]^2$$

Second moment about origin

Standard Deviation

$$S.D = \sqrt{Variance}$$

Gateway Classes : 7455 9612 84

Q.3 *If the probability density function of a random variable is given by*

AKTU 2021

$$f(x) = \begin{cases} k(1-x^2), & 0 \le x \le 1 \\ 0, & \text{elsewhere} \end{cases}$$

Find (i) k (ii) *Mean* (iii) *Varience* (iv) *Probability between* 0.1 *and* 0.2

(v) *Probability greater than* 0.5

Since $f(x)$ is a PDF

$\therefore$ (i) $f(x) \geq 0 \quad \forall x$

(ii) $\int_{-\infty}^{\infty} f(x)\,dx = 1$

From (ii)

$$\int_{-\infty}^{\infty} f(x)\,dx = 1$$

$$\int_{-\infty}^{0} f(x)\,dx + \int_{0}^{1} f(x)\,dx + \int_{1}^{\infty} f(x)\,dx = 1$$

$$0 + \int_0^1 k(1-x^2) + 0 = 1$$

$$k\left(x - \frac{x^3}{3}\right)_0^1 = 1$$

$$k\left[1 - \frac{1}{3}\right] = 1$$

$$\frac{2}{3}k = 1$$

$$k = \frac{3}{2}$$

$$k = 1.5$$

Gateway Classes : 7455 96 12 84

(ii) $\text{Mean} = \int_{-\infty}^{\infty} x f(x)\,dx$

$$E(x) = \int_{-\infty}^{0} x f(x)\,dx + \int_{0}^{1} x f(x)\,dx + \int_{1}^{\infty} x f(x)\,dx$$

$$E(x) = 0 + \int_{0}^{1} x K(1-x^2)\,dx + 0$$

$$E(x) = K\int_{0}^{1} (x - x^3)\,dx$$

$$E(x) = K\left[\frac{x^2}{2} - \frac{x^4}{4}\right]_0^1$$

$$E(x) = K\left[\frac{1}{2} - \frac{1}{4}\right]$$

$$E(x) = K \times \frac{1}{4}$$

$$E(x) = \frac{3}{2} \times \frac{1}{4}$$

$$\boxed{E(x) = \frac{3}{8}}$$

(iii) $E(x^2) = \int_{-\infty}^{\infty} x^2 f(x)\,dx$

$$E(x^2) = \int_{-\infty}^{0} x^2 f(x)\,dx + \int_{0}^{1} x^2 f(x)\,dx + \int_{1}^{\infty} x^2 f(x)\,dx$$

Gateway Classes 7455961284

$$E(x^2) = 0 + \int_0^1 x^2 k(1-x^2)dx + 0$$

$$E(x^2) = k\int_0^1 (x^2 - x^4)dx$$

$$E(x^2) = k\left(\frac{x^3}{3} - \frac{x^5}{5}\right)_0^1$$

$$E(x^2) = k\left(\frac{1}{3} - \frac{1}{5}\right]$$

$$E(x^2) = k\left(\frac{5-3}{15}\right)$$

$$E(x^2) = \frac{3}{2} \times \frac{2}{15} = \frac{1}{5}$$

$$\boxed{E(x^2) = \frac{1}{5}}$$

$$\text{variance} = E(x^2) - [E(x)]^2$$

$$= \frac{1}{5} - \left(\frac{3}{8}\right)^2 = \frac{1}{5} - \frac{9}{64}$$

$$= \frac{64 - 45}{320}$$

$$\boxed{\text{Var}(x) = \frac{19}{320} = 0.0594}$$

Gateway Classes : 7455 9612 84

(iv) $P(0.1 < x < 0.2) = \int_{0.1}^{0.2} f(x)\,dx$

$= \int_{0.1}^{0.2} k(1-x^2)\,dx$

$= k\left(x - \frac{x^3}{3}\right)_{0.1}^{0.2}$

$= \frac{3}{2}\left[0.2 - \frac{0.008}{3} - \left(0.1 - \frac{0.001}{3}\right)\right]$

$= \frac{3}{2}\left[0.1 - \frac{0.008}{3} + \frac{0.001}{3}\right]$

$P(0.1 < x < 0.2) = 0.1466$

(v) $P(0.5 < x < 1) = \int_{0.5}^{1} f(x)\,dx$

$= k\int_{0.5}^{1} (1-x^2)\,dx$

$= \frac{3}{2}\left[x - \frac{x^3}{3}\right]_{0.5}^{1}$

$= 0.3125$

Gateway Classes : 7455 961284

Q.4 *A continous random variable X has the probability density function*

$$f(x) = \begin{cases} kx & for\ 0 \le x < 2 \\ 2k & for\ 2 \le x < 4 \\ -kx + 6k & for\ 4 \le x \le 6 \\ 0 & elsewhere \end{cases}$$

Find (i)k (ii)Mean

Since $f(x)$ is PDF

$\therefore$ (i) $f(x) \ge 0\ \forall x$

(ii) $\int_{-\infty}^{\infty} f(x)\,dx = 1$

$$\int_{-\infty}^{\infty} f(x)\,dx = 1$$

$$\int_{-\infty}^{0} f(x)\,dx + \int_{0}^{2} f(x)\,dx + \int_{2}^{4} f(x)\,dx + \int_{4}^{6} f(x)\,dx + \int_{6}^{\infty} f(x)\,dx = 1$$

$$0 + k\int_{0}^{2} x\,dx + 2k\int_{2}^{4} dx + k\int_{4}^{6} (-x+6)\,dx + 0 = 1$$

Gateway Classes : 7455961284

$$k\left(\frac{x^2}{2}\right)_0^2 + 2k(x)_2^4 + k\left(-\frac{x^2}{2}+6x\right)_4^6 = 1$$

$$k(2-0) + 2k(4-2) + k\left[-18+36-(-8+24)\right] = 1$$

$$2k + 4k + k(18+8-24) = 1$$

$$2k + 4k + 2k = 1$$

$$8k = 1$$

$$\boxed{k = \frac{1}{8}}$$

(II) Mean

$$E(x) = \int_{-\infty}^{\infty} x\, f(x)\, dx$$

$$E(x) = \int_{-\infty}^{0} x f(x)\, dx + \int_{0}^{2} x f(x)\, dx + \int_{2}^{4} x f(x)\, dx + \int_{4}^{6} x f(x)\, dx + \int_{6}^{\infty} x f(x)\, dx$$

Gateway Classes : 7455961284

$$E(x) = k\int_0^2 x^2\,dx + 2k\int_2^4 x\,dx + k\int_4^6 (-x^2+6x)\,dx$$

$$= k\left(\frac{x^3}{3}\right)_0^2 + 2k\left(\frac{x^2}{2}\right)_2^4 + k\left(-\frac{x^3}{3}+\frac{6x^2}{2}\right)_4^6$$

$$= k\left[\frac{8}{3}-0+2(8-2)+\left(-72+3\times36+\frac{64}{3}-16\times3\right)\right]$$

$$= \frac{1}{8}\left[\frac{8}{3}+12+\left(-72+108+\frac{64}{3}-48\right)\right]$$

$$\boxed{E(x) = 3}$$

Gateway Classes : 7455 9612 84

Topic :*Random variables (PART-3)*

Gateway Classes : 7455 9612 84

Q. 1 Find the constant k so function $f(x)$ defined as follows be a density function:

$$\begin{cases} 1/k, & a \le x \le b \\ 0, & elsewhere \end{cases}$$

Q. 2 A continous random variable X has

$$\mathrm{f}(x) = \begin{cases} \frac{1}{2}(x+1), & for \ -1 \le x \le 1 \\ 0, & elsewhere \end{cases}$$

represent the density . Find mean, variance and standard deviation.

ENGINEERING MATHEMATICS

UNIT-4 : *Statistical Techniques – II*

Lec-04

Today's Target

- ***Binomial Probability Distribution***
- PYQ
- DPP

Gateway Classes : 7455 961284

Probability Distribution

Discrete probability distribution

(Based on Discrete random variable)

(i) **Binomial probability distribution** 2 L

(ii) **Poisson probability distribution** 2 L

Continous probability distribution

(Based on continous radom variable)

(i) **Normal probability Distribution** 2 L

Binomial probability Distribution

Bernoulli's Distribution

In Binomial distribution :

- Number of trials are finite (n)
- All trials are independent
- Each trial has only two outcomes (success or failure)

Gateway Classes : 7455 961 284

Binomial probability Distribution

Let a random variable X denote the number of successes in n independent trials in an experiment.

Let p be the probability of success and q be the probablity of failure in a single trial so that

$$p + q = 1$$

r successes can be obtained in n trials in ${}^{n}C_{r}$ ways.

Probability of r successes in n trials

$$P(X = r) = {}^{n}C_{r}\ \underbrace{P(S)\,P(S)\,\ldots\ldots\ldots P(S)}_{r \text{ times}}\ \underbrace{P(F)\,P(F)\,\ldots\ldots\ldots P(F)}_{(n-r) \text{ times}}$$

$$P(X = r) = {}^{n}C_{r}\ p\ p\ p\ \ldots\ldots\ldots p\ q\ q\ q\ \ldots\ldots\ldots q$$

$$\boxed{P(X = r) = {}^{n}C_{r}\ p^{r}\ q^{n-r}}$$

p = Prob. of success
q = Prob. of failure

n = No. of trials
r = No. of success

Gateway Classes : 7455 9612 84

Where $p + q = 1$

And $r = 0,\ 1,\ 2,\ 3\ \ldots$

$$P(X = 0) = n_{c_0}\, q^n$$

$$P(X = 1) = n_{c_1}\, p\, q^{n-1}$$

$$P(X = 2) = n_{c_2}\, p^2 q^{n-2}$$

$$\vdots \qquad\qquad \vdots$$

$$P(X = n) = n_{c_n}\, p^n$$

$n_{c_0}\, q^n$, $n_{c_1}\, p\, q^{n-1}$, $n_{c_2}\, p^2 q^{n-2}$ $\ldots\ldots\ldots\ldots$ $n_{c_n}\, p^n$ *are successive terms of binomial expansion* $(q + p)^n$

Gateway Classes : 7455 9612 84

➤ *If n independent trials constitute one experiment and this experiment is repeated N times.*

➤ *The frequuency of success is*

$$N\ n_{c_r}\ p^r\ q^{n-r}$$

And Binomial Distribution is $\boxed{N(q+p)^n}$

Mean, variance and standard deviation of Binomial distribution

(i) **Mean of Binomial Distribution,**

$$\boxed{\mu = np}$$

First moment about origin

(i) **Variance of Binomial Distribution,**

$$\boxed{\text{Variance} = npq}$$

Second moment about mean

(i) ***Standard Deviation,***

$$\sigma = \sqrt{npq}$$

GatewayClasses : 7455 9612 84

Recurrence or Recursion formula for Binomial Distribution

$$p(r+1) = \frac{n-r}{r+1} \frac{p}{q} p(r)$$

Q. 1 Comment on the following statement : For a Binomial distribution, mean is 6 and variance is 9.

Mean

$np = 6$ — ①

Variance

$npq = 9$ — ②

Divide ② by ①

$$\frac{npq}{np} = \frac{9}{6}$$

$$q = \frac{3}{2}$$

$q = 1.5$ (impossible)

Since, $0 \leq q \leq 1$

Given Statement is False

Gateway Classes : 7455 961284

Q. 2 *Ten coins are tossed simultaneously. Find the probalitity of getting at least seven heads.*

$P(H) = \frac{1}{2}$

$p = \frac{1}{2}$

$P(T) = \frac{1}{2}$

$q = \frac{1}{2}$

$n = 10$

$P(X=r) = {}^{n}C_{r} p^{r} q^{n-r}$

$P(r \geq 7) = P(r=7) + P(r=8) + P(r=9) + P(r=10)$

$= {}^{10}C_{7}\left(\frac{1}{2}\right)^{7} \times \left(\frac{1}{2}\right)^{3} + {}^{10}C_{8}\left(\frac{1}{2}\right)^{8}\left(\frac{1}{2}\right)^{2} + {}^{10}C_{9}\left(\frac{1}{2}\right)^{9}\left(\frac{1}{2}\right)^{1} + {}^{10}C_{10}\left(\frac{1}{2}\right)^{10}$

$= \left(\frac{1}{2}\right)^{10}\left[{}^{10}C_{7} + {}^{10}C_{8} + {}^{10}C_{9} + {}^{10}C_{10}\right]$

$= \left(\frac{1}{2}\right)^{10}\left[\frac{10 \times 9 \times 8}{3 \times 2} + \frac{10 \times 9}{2} + 10 + 1\right]$

$= \left(\frac{1}{2}\right)^{10} \times \left[120 + 45 + 10 + 1\right]$

$= 176 \times \left(\frac{1}{2}\right)^{10}$

$P(r \geq 7) = \frac{11}{64}$

Gateway Classes: 7455 96 2084

Q.3 *A die is thrown five times. If getting an odd number is a success, find the probability of getting at least four successes.*

$n = 5$

Getting an odd number is a success

$S = \{1, 2, 3, 4, 5, 6\}$

$P = \frac{3}{6} = \frac{1}{2}$

$P = \frac{1}{2}$ $\quad$ $q = \frac{1}{2}$

$P(x \geq 4) = P(x = 4) + P(x = 5)$

$= {}^5C_4 \left(\frac{1}{2}\right)^4 \times \frac{1}{2} + {}^5C_5 \left(\frac{1}{2}\right)^5$

$= \left(\frac{1}{2}\right)^5 \left[{}^5C_4 + {}^5C_5\right]$

$= \frac{1}{32}(5 + 1)$

$= \frac{1}{32} \times \frac{6}{16}^{3}$

$P(x \geq 4) = \frac{3}{16}$

Gateway Classes : 7455961284

Q.4 *A die is tossed thrice. A success is getting 1 or 6 on a toss. Find the mean and variance of the number of success.*

$S = \{1, 2, 3, 4, 5, 6\}$

Prob. of success

$$P = \frac{2}{6} = \frac{1}{3}$$

$$\boxed{P = \frac{1}{3}}$$

Prob. of failure

$$\boxed{q = \frac{2}{3}}$$

$n = 3$

Mean

$$\mu = np = 3 \times \frac{1}{3}$$

$$\boxed{\mu = 1}$$

Variance

$$Var(x) = npq = 3 \times \frac{1}{3} \times \frac{2}{3}$$

$$\boxed{Var(x) = \frac{2}{3}}$$

Gateway Classes 7455 9612 84

Q.5 *A binomial variable X satisfies the relation* $9P(X = 4) = P(X = 2)$ *when* $n = 6$.
Find the value of the parameter p and $P(X = 1)$.

$n = 6$

We know that

$$P(X = r) = {}^{n}C_{r}\, p^{r} q^{n-r}$$

$$P(X = 4) = {}^{6}C_{4}\, p^{4} q^{2}$$

$$P(X = 2) = {}^{6}C_{2}\, p^{2} q^{4}$$

According to question

$$9P(X = 4) = P(X = 2)$$

$$9 \times {}^{6}C_{4} \times p^{4} q^{2} = {}^{6}C_{2} \times p^{2} q^{4}$$

$$9 \times \frac{6!}{2! \times 4!} \times p^{2} = \frac{6!}{4! \times 2!} \times q^{2}$$

$$9p^{2} = q^{2}$$

$$9p^{2} = (1 - p)^{2}$$

Gateway Classes : 7455 961284

$$9p^2 = (1-p)^2$$

$$9p^2 = 1 + p^2 - 2p$$

$$8p^2 + 2p - 1 = 0$$

$$8p^2 + 4p - 2p - 1 = 0$$

$$4p(2p+1) - 1(2p+1) = 0$$

$$(4p-1)(2p+1) = 0$$

$$\boxed{p = \frac{1}{4}}$$

$p = -\frac{1}{2}$ (Rejected)

$$q = 1 - p$$

$$q = 1 - \frac{1}{4} = \frac{3}{4}$$

$$\boxed{q = \frac{3}{4}}$$

$$P(X=r) = {}^{n}C_{r}\, p^{r} q^{n-r}$$

$$P(X=1) = {}^{6}C_{1} \left(\frac{1}{4}\right)^{1} \times \left(\frac{3}{4}\right)^{5}$$

$$P(X=1) = {}^{3}\cancel{6} \times \frac{1}{\cancel{4}_{2}} \times \frac{3^5}{4^5}$$

$$P(X=1) = \frac{729}{2048}$$

$$\boxed{P(X=1) = 0.3559}$$

GatewayClasses: 7455 9612 84

Q.6 ***Fit a binomial distribution to the following frequency data :***

x:	0	1	2	3	4
f:	24	41	28	5	2

x	f	fx
0	24	0
1	41	41
2	28	56
3	5	15
4	2	8
	$\Sigma f = 100$	$\Sigma fx = 120$

$\Sigma f = N$

Mean

$$\mu = \frac{\Sigma fx}{\Sigma f}$$

$$= \frac{120}{100}$$

$$\mu = 1.2$$

For Binomial Distribution

$$\mu = np$$

$$1.2 = 4 \times p$$

$$p = 0.3$$

$$q = 0.7$$

Binomial Distribution

$$N(q+p)^n = 100(0.7+0.3)^4$$

Gateway Classes : 7455961284

Topic : *Binomial probability distribution*

Q.1 If the mean of a binomial distribution is 3 and the variabce is $\frac{3}{2}$, find the probability of obtaining at least 4 successes.

Q.2 The probability that a bomb dropped from a plane will strike the target is $\frac{1}{5}$.

If six bombs are dropped, find the probability that (i)exactly two will strike the target, (ii)at least two will trike the target.

Q.3 Fit a binimial distribution to the following frequency data:

x:	0	1	2	3	4
f:	30	62	46	10	2

GatewayClasses : 7455 9612 84

ENGINEERING MATHEMATICS

UNIT-4 : *Statistical Techniques – II*

Lec-05

Today's Target

- $\boldsymbol{Binomial\ probability\ distribution(Part-II)}$
- PYQ
- DPP

Gateway Classes : 7455 961284

Binomial probability Distribution

Let a random variable X denote the number of successes in n independent trials in an experiment.

Let p be the probability of success and q be the probablity of failure in a single trial so that

$$p + q = 1$$

r successes can be obtained in n trials in n_{C_r} ways.

Probability of r successes in n trials

$$P(X = r) = n_{C_r}\, P(S)\, P(S) \dots\dots\dots P(S)\quad P(F)\, P(F) \dots\dots\dots P(F)$$

$$P(X = r) = n_{C_r}\ \mathrm{p}\ \mathrm{p}\ \mathrm{p}\ \dots\dots\dots\ \mathrm{p}\ \mathrm{q}\ \mathrm{q}\ \mathrm{q}\ \dots\dots\dots\ \mathrm{q}$$

$$\boxed{P(X = r) = n_{C_r}\, p^r\, q^{n-r}}$$

Gateway Classes : 7455 9612 84

➤ *If n independent trials constitute one experiment and this experiment is repeated N times.*

➤ *The frequuency of success is*

$$N \; n_{c_r} \; p^r \; q^{n-r}$$

And Binomial Distrivution is $N(q+p)^n$

Mean, variance and standard deviation of Binomial distribution

(i) **Mean of Binomial Distribution,**

$$\mu = np$$

(i) **Variance of Binomial Distribution,**

$$\text{Variance} = npq$$

(i) ***Standard Deviation,***

$$\sigma = \sqrt{npq}$$

Third moment about Mean $= npq(q-p)$

Gateway Classes : 7455 9612 84

Q.1 *If 10% of bolts produced by a machine are defective, determine the probability that out of 10 bolts choosen at random. (i)1 (ii)None (iii)at most 2 bolts will be defective.*

$$P(\text{defective}) = \frac{10}{100} = \frac{1}{10}$$

$$P = \frac{1}{10}$$

$$P(\text{Non Defective}) = 1 - \frac{1}{10}$$

$$q = \frac{9}{10}$$

$$n = 10$$

(i) $P(X = r) = {}^{n}C_{r}\, p^{r} q^{n-r}$

$$P(X = 1) = {}^{10}C_{1} \left(\frac{1}{10}\right)^{1} \times \left(\frac{9}{10}\right)^{9}$$

$$P(X = 1) = 10 \times \frac{1}{10} \times (0.9)^{9}$$

$$P(X = 1) = 0.3874$$

Gateway Classes : 7455 961284

(ii) $P(X=r) = {}^{n}C_r\, p^r q^{n-r}$

$$P(X=0) = {}^{10}C_0 \left(\frac{1}{10}\right)^0 \times \left(\frac{9}{10}\right)^{10}$$

$$P(X=0) = 1 \times 1 \times (0.9)^{10}$$

$$\boxed{P(X=0) = 0.3487}$$

(iii) $P(r \le 2) = P(X=0) + P(X=1) + P(X=2)$ — ①

$$P(X=2) = {}^{10}C_2\, p^2 q^8$$

$$P(X=2) = \frac{10!}{8! \times 2!} \times \left(\frac{1}{10}\right)^2 \times \left(\frac{9}{10}\right)^8$$

$$= \frac{10 \times 9 \times 8!}{8! \times 2} \times \frac{1}{100} \times (0.9)^8$$

$$\boxed{P(X=2) = 0.1937}$$

Required Probability

$$P(r \le 2) = 0.3487 + 0.3874 + 0.1937$$

$$\boxed{P(r \le 2) = 0.9298}$$

Gateway Classes : 7455 961284

Q.2 *If the probability of hitting a target is 10% and 10 shots are fired independently. What is the probability that the target will be hit at least once?* **AKTU 2019**

$P(\text{Hitting a target}) = \frac{10}{100}$

$P = \frac{1}{10}$

$P(\text{Not Hitting target}) = 1 - \frac{1}{10}$

$q = \frac{9}{10}$

$n = 10$

$P(r \geq 1) = 1 - P(x = 0)$

$= 1 - {}^{n}C_{r} \, p^{r} q^{n-r}$

$= 1 - {}^{10}C_{0} \left(\frac{1}{10}\right)^{0} \times \left(\frac{9}{10}\right)^{10}$

$= 1 - 1 \times 1 \times (0.9)^{10}$

$P(r \geq 1) = 0.6513$

Gateway Classes : 7455 961284

Q.3 *In 800 families with 5 children each, how many families would be expected to have (i) 3 boys and 2 girls, (ii) 2 boys and 3 girls, (iii) no girl (iv) at the most two girls. Assume probabilities for boys and girls to be equal.* **AKTU-2017** V.imp

$N = 800$

$n = 5$

$P(\text{Having a girl}) = \frac{1}{2}$

$$p = \frac{1}{2}$$

$P(\text{Having a boy}) = \frac{1}{2}$

$$q = \frac{1}{2}$$

(i) Expected number of families having 3 boys and 2 girls

$$N\,P(X = 2) = N\,{}^{n}C_{r}\,p^{r}\,q^{n-r}$$

$$= 800 \times {}^{5}C_{2}\left(\frac{1}{2}\right)^{2} \times \left(\frac{1}{2}\right)^{3}$$

$$= 800 \times \frac{5!}{3!2!} \times \left(\frac{1}{2}\right)^{5}$$

$$= 800 \times \frac{5 \times 4 \times 3!}{3! \times 2} \times \frac{1}{32}$$

Gateway Classes: 7455961284

$$N(X=2) = 250$$

(ii) Number of families having 2 boys and 3 Girls

$$NP(X=3) = N\,{}^{n}C_{\gamma}\, p^{\gamma} q^{n-\gamma}$$

$$= 800 \times {}^{5}C_{3} \left(\frac{1}{2}\right)^{3} \times \left(\frac{1}{2}\right)^{2}$$

$$= 800 \times \frac{5!}{3! \times 2!} \times \left(\frac{1}{2}\right)^{5}$$

$$NP(X=3) = 250$$

(iii) Number of families having No girl

$$NP(X=0) = N\,{}^{n}C_{\gamma}\, p^{\gamma} q^{n-\gamma}$$

$$= 800 \times {}^{5}C_{0} \left(\frac{1}{2}\right)^{0} \times \left(\frac{1}{2}\right)^{5}$$

$$= \frac{800}{32} \times 1 \times \frac{1}{32}$$

$$NP(X=0) = 25$$

Gateway Classes : 7455 961284

(iv) Number of families having atmost two Girls

$$NP(r \le 2) = N[P(x=0) + P(x=1) + P(x=2)]$$

$$P(x=1) = {}^{5}C_1 \, p^r \, q^{n-r}$$

$$= 5 \times \left(\frac{1}{2}\right)^1 \times \left(\frac{1}{2}\right)^4$$

$$= 5 \times \frac{1}{25}$$

$$= \frac{5}{32}$$

$$NP(r \le 2) = NP(x=0) + NP(x=1) + NP(x=2)$$

$$= 25 + \frac{25}{\cancel{800}} \times \frac{5}{\cancel{32}} + 250$$

$$= 25 + 125 + 250$$

$$NP(r \le 2) = 400$$

Gateway Classes : 7455 961284

Q. 4 *A student is given a true false examination with 8 questions. If he correct at least 7 questions, he passes the examination. Find the probability that he will pass given that he guesses all questions.*

$n = 8$

$p = \frac{1}{2}$

$q = \frac{1}{2}$

Prob. that he will pass

$P(r \geq 7) = P(x=7) + P(x=8)$

$= {}^8C_7 \left(\frac{1}{2}\right)^7 \times \left(\frac{1}{2}\right) + {}^8C_8 \left(\frac{1}{2}\right)^8 \left(\frac{1}{2}\right)^0$

$P(r \geq 7) = 8 \times \left(\frac{1}{2}\right)^8 + 1 \times \left(\frac{1}{2}\right)^8$

$= \left(\frac{1}{2}\right)^8 (8+1)$

$= \frac{9}{2^8}$

$P(r \geq 7) = 0.035$

Gateway Classes : 7455 961284

Q.5 *Six dies are thrown 729 times. How many times do you expect at least three dice to show a five or six?*

$N = 729$

$n = 6$

$S = \{1, 2, 3, 4, 5, 6\}$

$P(\text{having } 5 \text{ or } 6) = \frac{2}{6} = 3$

$$p = \frac{1}{3}$$

$P(\text{having Not } 5 \text{ or } 6) = \frac{4}{6} = \frac{2}{3}$

$$q = \frac{2}{3}$$

$$NP(x \geq 3) = N[P(x=3) + P(x=4) + P(x=5) + P(x=6)]$$

$$= 729\left[{}^6C_3\left(\frac{1}{3}\right)^3\left(\frac{2}{3}\right)^3 + {}^6C_4\left(\frac{1}{3}\right)^4\left(\frac{2}{3}\right)^2 + {}^6C_5\left(\frac{1}{3}\right)^5\left(\frac{2}{3}\right)^1 + {}^6C_6\left(\frac{1}{3}\right)^6\left(\frac{2}{3}\right)^0\right]$$

$$NP(x \geq 3) = 729\left[{}^6C_3 \times \frac{8}{3^6} + {}^6C_4 \times \frac{4}{3^6} + {}^6C_5 \times \frac{2}{3^6} + {}^6C_6 \times \frac{1}{3^6}\right]$$

Gateway Classes 7455 961284

$$NP(r \geq 3) = \frac{729}{3^6}\left[8\,{}^6C_3 + 4\,{}^6C_4 + 2\,{}^6C_5 + {}^6C_6\right]$$

$$= \frac{729}{729}\left[8 \times \frac{6!}{3! \times 3!} + 4 \times \frac{6!}{4! \times 2!} + 2 \times 6 + 1\right]$$

$$= \frac{8 \times 6 \times 5 \times 4 \times 3!}{3 \times 2 \times 1 \times 3!} + \frac{4 \times 6 \times 5 \times 4!}{4! \times 2 \times 1} + 12 + 1\Big]$$

$$= 160 + 60 + 12 + 1$$

$$\boxed{NP(r \geq 3) = 233}$$

GW GATEWAY CLASSES
Gateway Classes 7455961284

Topic : Binomial probability distribution(Part – II)

Q.1 Out of 800 families with 4 children each, how many families would be expected to have (i)2 boys and 2 girls (ii)at least one boy (iii)no girl (iv)almost two girls? Assume equal probabilities for boys and girls. UPTU 2014

Q.2 A bag contains 5 white , 7 red and 8 black balls. If four balls are drawn, one by one, with replacement. what is the probability that

(i) none is white

(ii) all are white

(iii) at least on is white

(iv) only 2 are white ?

Q.3 Five cards are drawn successively with repelacement from a wellshuffled deck of 52 cards. What is the probability that

(i) All the five cards are spades

(ii) only three are spades ?

(iii) none is spade ?

Gateway Classes : 7455 961284

ENGINEERING MATHEMATICS

UNIT-4 : *Statistical Techniques – II*

Lec-06

Today's Target

- $\boldsymbol{Poisson\ \ Probability\ Distribution(Part-I)}$
- PYQs
- DPPs

Gateway Classes : 7455 961284

Poisson Probability Distribution

- Poisson Distribution is a limiting case of binomial distribution

$$P(X = r) = \frac{\lambda^r e^{-\lambda}}{r!} \qquad (r = 0, 1, 2, 3, \dots)$$

- Recurrence formula for the Poisson Distribution

$$P(r+1) = \frac{\lambda}{(r+1)} P(r) \qquad (r = 0, 1, 2, 3, \dots.)$$

- Mean and Variance of the Poisson Distribution

$$Mean = \lambda$$

$$Variance = \lambda$$

Gateway Classes : 7455 9612 84

Important Derivations:

1. Show that Poisson Distribution is a limiting case of binomial distribution (A.K.T.U. 2021)

In Binomial Distribution

n = number of trials

P = Prob. of success

- If n is very large and p is very small then Binomial distribution become complicated
- If we assume that $n \to \infty$ and $p \to 0$ such that $np = \lambda$ is finite

Then we get poisson approximation to binomial distribution.

$$P(X = r) = \frac{e^{-\lambda} \lambda^r}{r!}$$

Proof :

$$P(X = r) = {}^{n}C_r \, p^r q^{n-r}$$

Gatway Classes. 7455961284

$$P(X=r) = {}^{n}C_r \, p^r (1-p)^{n-r} \quad \{p+q=1\}$$

$$P(X=r) = \frac{n!}{(n-r)!\,r!} p^r \times (1-p)^n (1-p)^{-r}$$

$$P(X=r) = \frac{n!}{(n-r)!\,r!} \times \left(\frac{\lambda}{n}\right)^r \times \frac{\left(1-\frac{\lambda}{n}\right)^n}{\left(1-\frac{\lambda}{n}\right)^r}$$

$$P(X=r) = \frac{\lambda^r \times n(n-1)(n-2)\cdots\cdots(n-r+1)\,(n-r)!\,\left(1-\frac{\lambda}{n}\right)^n}{r!\times (n-r)!\times n^r \times \left(1-\frac{\lambda}{n}\right)^r}$$

$$P(X=r) = \frac{\lambda^r}{r!} \times \frac{n}{n} \times \left(\frac{n-1}{n}\right)\left(\frac{n-2}{n}\right)\cdots\cdots\left(\frac{n-r+1}{n}\right) \frac{\left(1-\frac{\lambda}{n}\right)^n}{\left(1-\frac{\lambda}{n}\right)^r}$$

$$P(X=r) = \frac{\lambda^r}{r!} \times 1 \times \left(1-\frac{1}{n}\right)\left(1-\frac{2}{n}\right)\cdots\cdots\left(1-\frac{r}{n}+\frac{1}{n}\right) \frac{\left(1-\frac{\lambda}{n}\right)^n}{\left(1-\frac{\lambda}{n}\right)^r}$$

Gateway Classes : 7455 9612 84

$$P(X=r) = \lim_{n\to\infty} \frac{\lambda^r}{r!}\left(1-\frac{1}{n}\right)\left(1-\frac{2}{n}\right)\cdots\cdots\left(1-\frac{r}{n}+\frac{1}{n}\right)\frac{\left(1-\frac{\lambda}{n}\right)^n}{\left(1-\frac{\lambda}{n}\right)^r}$$

$$= \frac{\lambda^r}{r!}(1-0)(1-0)\cdots\cdots(1-0+0)\frac{\lim_{n\to\infty}\left(1-\frac{\lambda}{n}\right)^n}{\lim_{n\to\infty}\left(1-\frac{\lambda}{n}\right)^r}$$

$$= \frac{\lambda^r}{r!} \times 1 \times 1 \cdots\cdots \times 1 \times \frac{e^{-\lambda}}{1}$$

$$= \frac{\lambda^r}{r!} \times e^{-\lambda}$$

$$\boxed{P(X=r) = \frac{\lambda^r e^{-\lambda}}{r!}}$$ Poisson Distribution

$$\lim_{n\to\infty}\left(1-\frac{x}{n}\right)^n = e^{-x}$$

$$\lim_{n\to\infty}\left(1-\frac{x}{n}\right)^r = 1$$

Gateway Classes : 7455 9612 84

Poisson Distribution

$$P(X = r) = \frac{\lambda^r e^{-\lambda}}{r!} \qquad (r = 0, 1, 2, 3, \dots)$$

- λ *is called the parameter of the distribution.*

λ = np (Finite)

- *The sum of the Probability* $P(r)$ *for* $r = 0, 1, 2, 3, \dots \dots$ *is* 1

$$\sum P(r) = 1$$

P(X=0) + P(X=1) + P(X=2) + P(X=3) . . . . . = 1

Gateway Classes: 7455 9612 84

2. Recurrence formula for the Poisson Distribution

By poisson Distribution

$$P(r) = \frac{\lambda^r e^{-\lambda}}{r!} \quad \text{——} \quad (1)$$

Replace r by $r+1$

$$P(r+1) = \frac{\lambda^{r+1} e^{-\lambda}}{(r+1)!}$$

$$P(r+1) = \frac{\lambda^r \times \lambda^1 \times e^{-\lambda}}{(r+1) \times r!} \quad \text{——} \quad (2)$$

Divide (2) by (1)

$$\frac{P(r+1)}{P(r)} = \frac{\lambda^r \times \lambda \times e^{-\lambda}}{(r+1) \times r!} \times \frac{r!}{\lambda^r \times e^{-\lambda}}$$

$$\frac{P(r+1)}{P(r)} = \frac{\lambda}{r+1}$$

$$\boxed{P(r+1) = \frac{\lambda}{r+1} P(r)}$$

Gateway Classes : 7455 961284

3. Mean and Variance of the Poisson Distribution

By Poisson Distribution

$$P(r) = \frac{e^{-\lambda}\lambda^{r}}{r!}$$

Mean

$$\mu = E(r) = \sum_{r=0}^{\infty} r\,p(r)$$

$$\mu = \sum_{r=0}^{\infty} \frac{r\,e^{-\lambda}\lambda^{r}}{r!}$$

$$\mu = e^{-\lambda} \sum_{r=1}^{\infty} \frac{r\,\lambda^{r}}{r(r-1)!}$$

$$\mu = e^{-\lambda} \sum_{r=1}^{\infty} \frac{\lambda^{r}}{(r-1)!}$$

$$\mu = e^{-\lambda}\left(\frac{\lambda}{0!} + \frac{\lambda^2}{1!} + \frac{\lambda^3}{2!} + \frac{\lambda^4}{3!} + \cdots\right)$$

$$\mu = e^{-\lambda}\lambda\left(1 + \frac{\lambda}{1!} + \frac{\lambda^2}{2!} + \frac{\lambda^3}{3!} + \cdots\right)$$

$$\text{Mean} = e^{-\lambda} \times \lambda \times e^{\lambda} = \lambda \times e^{0}$$

$$\boxed{\text{Mean} = \lambda}$$

Gateway Classes: 7455961284

Variance

$$\sigma^2 = E(x^2) - [E(x)]^2$$

$$\sigma^2 = \sum_{x=0}^{\infty} x^2 P(x) - (\lambda)^2$$

$$\sigma^2 = \sum_{x=0}^{\infty} \frac{x^2 e^{-\lambda} \lambda^x}{x!} - \lambda^2$$

$$\sigma^2 = e^{-\lambda} \sum_{x=1}^{\infty} \frac{x^2 \lambda^x}{x!} - \lambda^2$$

$$\sigma^2 = e^{-\lambda}\left(\frac{1^2 \times \lambda}{1!} + \frac{2^2 \times \lambda^2}{2!} + \frac{3^2 \times \lambda^3}{3!} + \dots\right) - \lambda^2$$

$$\sigma^2 = e^{-\lambda} \times \lambda\left(1 + \frac{2\lambda}{1!} + \frac{3\lambda^2}{2!} + \frac{4\lambda^3}{3!} + \dots\right) - \lambda^2$$

$$\sigma^2 = \lambda e^{-\lambda}\left[1 + \frac{\lambda+\lambda}{1!} + \frac{\lambda^2+2\lambda^2}{2!} + \frac{\lambda^3+3\lambda^3}{3!} + \dots\right) - \lambda^2$$

$$\sigma^2 = \lambda e^{-\lambda}\left[1 + \frac{\lambda}{1!} + \frac{\lambda}{1!} + \frac{\lambda^2}{2!} + \frac{2\lambda^2}{2!} + \frac{\lambda^3}{3!} + \frac{3\lambda^3}{3!} + \dots\right) - \lambda^2$$

$$\sigma^2 = \lambda e^{-\lambda}\left[\left(1 + \frac{\lambda}{1!} + \frac{\lambda^2}{2!} + \dots\right) + \left(\frac{\lambda}{1!} + \frac{2\lambda^2}{2!} + \frac{3\lambda^3}{3!} + \dots\right)\right) - \lambda^2$$

Gateway Classes : 7455 961284

$$\sigma^2 = \lambda e^{-\lambda}\left[\left(1 + \frac{\lambda}{1!} + \frac{\lambda^2}{2!} + \frac{\lambda^3}{3!} + \cdots\right) + \lambda\left(1 + \frac{2\lambda}{2!} + \frac{3\lambda^2}{3!} + \frac{4\lambda^3}{4!} + \cdots\right) - \lambda^2\right.$$

$$\sigma^2 = \lambda e^{-\lambda}\left[e^{\lambda} + \lambda\left(1 + \frac{\lambda}{1!} + \frac{\lambda^2}{2!} + \frac{\lambda^3}{3!} + \cdots\right) - \lambda^2\right.$$

$$\sigma^2 = \lambda e^{-\lambda}\left[e^{\lambda} + \lambda e^{\lambda}\right] - \lambda^2$$

$$\sigma^2 = \lambda e^{-\lambda} e^{\lambda} + \lambda^2 e^{-\lambda} e^{\lambda} - \lambda^2$$

$$\sigma^2 = \lambda + \lambda^2 - \lambda^2$$

$$\sigma^2 = \lambda$$

Variance $= \lambda$

Gateway Classes : 7455 9612 84

Q.1 *If a random variable X follow a poisson distribution such that* $P(X=2)=9P(X=4)+90P(X=6)$. *Find mean and variance of X.* **(A.K.T.U. 2023)**

For poisson Distribution

$$P(X=x)=\frac{e^{-\lambda}\lambda^{x}}{x!}$$

$$P(X=2)=9P(X=4)+90P(X=6)$$

$$\frac{e^{-\lambda}\lambda^{2}}{2!}=9\times\frac{e^{-\lambda}\lambda^{4}}{4!}+\frac{90\times e^{-\lambda}\lambda^{6}}{6!}$$

$$\frac{e^{-\lambda}\lambda^{2}}{2!}=e^{-\lambda}\left(\frac{9\lambda^{4}}{4!}+\frac{90\lambda^{6}}{6!}\right)$$

$$\frac{\lambda^{2}}{2}=\lambda^{2}\left(\frac{3\times 3\lambda^{2}}{4\times 3\times 2\times 1}+\frac{30\times 3\lambda^{4}}{6\times 5\times 4\times 3\times 2\times 1}\right)$$

$$\frac{1}{2}=\frac{3\lambda^{2}}{8}+\frac{\lambda^{4}}{8}$$

$$\frac{1}{2}=\frac{3\lambda^{2}+\lambda^{4}}{8}$$

$$4=3\lambda^{2}+\lambda^{4}$$

$$\lambda^{4}+3\lambda^{2}-4=0$$

Gateway Classes : 7455961284

Put $\lambda^2 = x$

$$x^2 + 3x - 4 = 0$$

$$x^2 + 4x - x - 4 = 0$$

$$x(x+4) - 1(x+4) = 0$$

$$(x-1)(x+4) = 0$$

$x = 1$	$x = -4$
$\lambda^2 = 1$	$\lambda^2 = -4$
$\lambda = \pm 1$	(Reject)
$\lambda = 1$	

Hence

$$\lambda = 1$$

Mean = 1

Variance = 1

Gateway Classes 7455 9612 84

Q.2 *If the variance of the Poisson distribution is 2, find the probabilities for* $r = 1, 2, 3, 4$ *from the recurrence relation of the Poisson distribution. Also, find* $P(r \geq 4)$.

(M.T.U. 2013)

Given

$\lambda = 2$

By poisson Distribution

$$P(r) = \frac{e^{-\lambda} \lambda^r}{r!}$$

$$P(0) = \frac{e^{-2} \times \lambda^0}{0!}$$

$$P(0) = e^{-2}$$

$$P(0) = 0.1353$$

By Recurrence Relation

$$P(r+1) = \frac{\lambda}{r+1} P(r)$$

$$P(0+1) = \frac{2}{0+1} P(0)$$

$$P(1) = 2\, P(0)$$

$$\boxed{P(1) = 0.2706}$$

Gateway Classes : 7455 961284

$$P(1+1) = \frac{2}{1+1} P(1)$$

$$P(2) = P(1)$$

$$\boxed{P(2) = 0.2706}$$

$$P(2+1) = \frac{2}{2+1} P(2)$$

$$P(3) = \frac{2}{3} \times 0.2706$$

$$\boxed{P(3) = 0.1804}$$

$$P(3+1) = \frac{2}{3+1} P(3)$$

$$P(4) = \frac{1}{2} \times 0.1804$$

$$\boxed{P(4) = 0.0902}$$

$$P(x \geq 4) = 1 - P(x < 4)$$

$$P(x \geq 4) = 1 - [P(0) + P(1) + P(2) + P(3)]$$

$$P(x \geq 4) = 1 - 0.8569$$

$$\boxed{P(x \geq 4) = 0.1431}$$

Gateway Classes : 7455 9612 84

Q.3 *Using Poisson distribution, find the probability that the ace of spades will be drawn from a pack of well – shuffled cards at least once in 104 consecutive trials.* **(U.P.T.U. 2015)**

$P(\text{ace of spades}) = \frac{1}{52}$

$p = \frac{1}{52}$

$n = 104$

$\lambda = np$

$\lambda = 104 \times \frac{1}{52}$

$\lambda = 2$

By Poisson Distribution

$$P(r) = \frac{e^{-\lambda} \lambda^r}{r!}$$

$$P(r \geq 1) = 1 - P(x = 0)$$

$$= 1 - \frac{e^{-2} \times \lambda^0}{0!} = 1 - e^{-2}$$

$$P(r \geq 1) = 0.8647$$

Gateway Classes 7455 961284

DPP- 06

Topic :*Poisson Probability Distribution (Part – I)*

Q.1 In a poisson distribution, $P(r)$ for $r = 0$ is 10%, find the mean.

Q.2 If X poisson variate such that $P(X = 1) = P(X = 2)$. Find (i) Mean (ii) $P(X = 4)$

Gateway Classes : 7455 961284

ENGINEERING MATHEMATICS

UNIT-4 : *Statistical Techniques – II*

Lec-07

Today's Target

- $\boldsymbol{Poisson\ Probability\ Distribution(Part-II)}$
- PYQs
- DPPs

Gateway Classes : 7455 961284

Poisson Probability Distribution

- **Poisson Distribution is a limiting case of binomial distribution**

$$P(X = r) = \frac{\lambda^r e^{-\lambda}}{r!} \qquad (r = 0, 1, 2, 3, \dots)$$

- **Recurrence formula for the Poisson Distribution**

$$P(r+1) = \frac{\lambda}{(r+1)} P(r) \qquad (r = 0, 1, 2, 3, \dots.)$$

- **Mean and Variance of the Poisson Distribution**

$$\text{Mean} = \lambda$$

$$\text{Variance} = \lambda$$

$$\lambda = np$$

Gateway Classes : 7455 9612 84

Q.1 *It is given that 2% of the electric bulbs manufactured by a company are defective. Using Poisson distribution, find the probability that a sample of 200 bulbs will contain*

(i) no defective bulb (ii) two defective bulbs (iii) at the most three defective bulbs. (**G.B.T.U. 2011**)

Prob. of Defective bulbs

$$P = \frac{2}{100}$$

$$P = 0.02$$

$$n = 200$$

$$\lambda = 0.02 \times 200$$

$$\lambda = 4$$

By poisson distribution

$$P(X=r) = \frac{e^{-\lambda} \lambda^r}{r!}$$

(i) Prob. of no defective bulb ($r=0$)

$$P(X=0) = \frac{e^{-4} \times (4)^0}{0!} = e^{-4}$$

$$P(X=0) = 0.018315$$

(ii) Prob. of two defective ($r=2$)

$$P(X=2) = \frac{e^{-4}(4)^2}{2!} = \frac{16}{2} e^{-4}$$

$$= 8 e^{-4}$$

Gateway Classes : 7455 961 204

$$P(X=2) = 8e^{-4}$$

$$P(X=2) = 0.146525$$

(iii) Prob. of atmost 3 defective bulbs $(x \leq 3)$

$$P(x \leq 3) = P(X=0) + P(X=1) + P(X=2) + P(X=3)$$

$$= e^{-4} + \frac{e^{-4} \times 4}{1!} + \frac{8e^{-4}}{2!} + \frac{e^{-4}(4)^3}{3!}$$

$$= e^{-4} + 4e^{-4} + 8e^{-4} + \frac{32}{3}e^{-4}$$

$$P(x \leq 3) = e^{-4}\left(13 + \frac{32}{3}\right)$$

$$= e^{-4}\left(\frac{39+32}{3}\right)$$

$$= \frac{71}{3}e^{-4}$$

$$\boxed{P(x \leq 3) = 0.43347}$$

Gateway Classes : 7455 9612 84

Q.2 *A manufacturer knows that the condensors he makes contain on an average 1% of defectives. He packes them in boxes of 100. What is the probability that a box picked at random will contain 4 or more faulty condensors?*

Prob. of defective condensors

$$P = \frac{1}{100}$$

$$\boxed{P = 0.01}$$

$$n = 100$$

$$\lambda = np = 100 \times 0.01$$

$$\boxed{\lambda = 1}$$

By poisson distribution

$$P(X = r) = \frac{e^{-\lambda} \lambda^{r}}{r!}$$

Prob. of 4 or more faulty condensor

$(r \geq 4)$

$$P(r \geq 4) = 1 - P(r < 4)$$

Gateway Classes : 7455961284

$$P(x \geq 4) = 1 - [P(x=0) + P(x=1) + P(x=2) + P(x=3)]$$

$$= 1 - \left[\frac{e^{-1}(1)^0}{0!} + \frac{e^{-1} \times (1)^1}{1!} + \frac{e^{-1} \times 1^2}{2!} + \frac{e^{-1} \times 1^3}{3!}\right]$$

$$= 1 - \left[e^{-1} + e^{-1} + \frac{1}{2}e^{-1} + \frac{1}{6}e^{-1}\right]$$

$$= 1 - e^{-1}\left(2e^{-1} + \frac{e^{-1}}{2} + \frac{e^{-1}}{6}\right)$$

$$= 1 - e^{-1}\left(\frac{12e^{-1} + 3e^{-1} + e^{-1}}{6}\right)$$

$$= \left(1 - \frac{8}{3}e^{-1}\right) = 0.018988$$

$$P(x \geq 4) = 0.018988$$

Gateway Classes : 7455961284

Q.3 *In a certain factory turning out razor blades, there is a small chance of* 0.002 *for any blade to be defective. The blades are supplied in packets of* 10*. Calculate the approximate number of packets containing no defective, one defective and two defective blades in a consignment of* $10,000$ *packets.* (*Given:* $e^{-0.02} = 0.9802$) (*A.K.T.U. 2018*)

Prob. of defective blade

$$P = 0.002$$

$$n = 10$$

$$N = 10,000$$

$$\lambda = nP = 0.002 \times 10$$

$$\boxed{\lambda = 0.02}$$

By Poisson Distribution

$$P(X = r) = \frac{e^{-\lambda} \lambda^r}{r!}$$

(i) Number of packets containing no defective blades

$$P(X = 0) = \frac{e^{-0.02} \times (0.02)^0}{0!} = e^{-0.02}$$

Gateway Classes : 7455 96 12 84

$P(X=0) = e^{-0.02}$

$P(X=0) = 0.9802$

Number of packets

$= 10,000 \times 0.9802$

$= 9802$

(ii) $P(X=1) = \dfrac{e^{-0.02} \times (0.02)^1}{1!}$

$P(X=1) = 0.0196$

Number of packets having 1 defective blades

$= 0.0196 \times 10000$

$= 196$

(iii) $P(X=2) = \dfrac{e^{-0.02} \times (0.02)^2}{2!}$

$= 0.000196$

Number of packets having 2 defective blades

$= 0.000196 \times 10,000 = 1.96$

$= 2$

Gateway Classes : 7455 961284

Q.4 *The number of accidents in a year involving taxi drivers in a city follows a Poisson distribution with mean equal to 3. Out of 1000 taxi drivers, find approximately the numberof drivers with (i)no accidents (ii)more than 3 accidents in a year.* **(A.K.T.U. 2021)**

Given

$\lambda = 3$

$N = 1000$

By poisson Distribution

$$P(X=r) = \frac{e^{-\lambda}\,\lambda^{r}}{r!}$$

(i) $$P(X=0) = \frac{e^{-3}\,(3)^{0}}{0!} = e^{-3}$$

$$P(X=0) = e^{-3} = 0.04978$$

Number of drivers having no accidents

$$= 0.04978 \times 1000$$

$$= 49.7$$

$$= 50$$

Gateway Classes : 7455 9612 84

(11) $P(x>3) = 1-[P(x=0)+P(x=1)+P(x=2)+P(x=3)]$

$$= 1-\left[e^{-3}+\frac{e^{-3}\times(3)^1}{1!}+\frac{e^{-3}\times(3)^2}{2!}+\frac{e^{-3}\times(3)^3}{3!}\right]$$

$$= 1-\left[e^{-3}+3e^{-3}+\frac{9}{2}e^{-3}+\frac{9}{2}e^{-3}\right]$$

$$= 1-e^{-3}(4+4.5+4.5)$$

$$= 1-13e^{-3} = 0.3527$$

Number of drivers having more than 3 accidents

$$= 0.3527\times 1000 = 352.7$$

$$= 353$$

GW GATEWAY CLASSES
Gateway Classes : 7455961284

Q.5 *The following table gives the no. of days in a 50 day period during which automobile accidents occured in a city.*

No. of accidents :	0	1	2	3	4
No. of days (f) :	21	18	7	3	1

Fit a Poisson distribution to the data. **(A.K.T.U. 2022)**

x	f	fx	Theoritical frequencies
0	21	0	20
1	18	18	18
2	7	14	8
3	3	9	3
4	1	4	1
	$\Sigma f = 50$	$\Sigma fx = 45$	

$$\text{Mean} = \frac{\Sigma fx}{\Sigma f}$$

$$\lambda = \frac{45}{50}$$

$$\boxed{\lambda = 0.9}$$

$$N = 50$$

Gateway Classes : 7455 9612 84

Poisson Distribution $= N \frac{e^{-\lambda} \lambda^r}{r!}$

$$= \frac{50 \times e^{-0.9} (0.9)^r}{r!}$$

Wher $r = 0$

$= 20.32 = 20$

When $r = 0$

$= 18.29 = 18$

When $r = 2$

$= 8.2 = 8$

When $r = 3$

$= 2.46$

$= 3$

When $r = 4$

$= 0.555$

$= 1$

Gateway Classes : 7455 9612 84

DPP- 07

Topic : *Poisson Probability Distribution (Part – II)*

Q. 1 In a certain factory manufacturing razor blades, there is a small chance of 0.002 for any blade to be defective. The blades are supplied in packets of 10. Use suitable distribution to calculate the approximate number of packets containing no defective, one defective and two defective blades respectively in a consignment of 50,000 packets. (*AKTU 2018*)

Q. 2 The distribution of the number of road accidents per day in a city is Poisson with mean 4. Find the number of days out of 100 days when there will be (i) no accident (ii) at least 2 accidents (iii) at most 3 accidents (iv) between 2 and 5 accidents.

Q. 3 Fit a Poisson distribution to the following data and calculate theoretical frequencies.

Deaths:	0	1	2	3	4
Frequencies:	122	60	15	2	1

Q. 4 Fit a Poisson distribution to the following data given the number of yeast cells per square for 400 squares:

No. of cells per sq. :	0	1	2	3	4	5	6	7	8	9	10
No. of squares :	103	143	98	42	8	4	2	0	0	0	0

Gateway Classes : 7455961284

ENGINEERING MATHEMATICS

UNIT-4 : *Statistical Techniques – II*

Lec-08

Today's Target

- $Normal\ Probability\ Distribution (Part - I)$
- PYQs
- DPPs

Gateway Classes : 7455 961284

Normal Distribnution

- The normal distribution is a continous distribution based on continous random variable
- A continous random variable X is said to have a normal distribution with parameters $\mu(Mean)$ and $\sigma^2(Variance)$ if the probability density function is given by

$$f(x) = \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{1}{2}\left(\frac{x-\mu}{\sigma}\right)^2}$$

(i) $f(x) \geq 0$

(ii) $\int_{-\infty}^{\infty} f(x)dx = 1.$

$-\infty < x < \infty$


μ = Mean

σ = Standard deviation

Gateway Classes : 7455 961284

Normal Curve

- The graph of normal *Distribnution is called normal curve.*
- *It is bell shaped and symmetrical about Mean μ.*



- *In normal distribution Mean, median and mode coincide.*
- *Total area under the curve above x − axis is 1*

Gateway Classes : 7455 9612 84

Standard Normal Distribution

- *If X is a Normal random variable with mean μ and standard deviation σ, then a random Variable Z with Mean = 0 and standard deviation = 1 is called standard normal random variable.*

$$Z = \frac{x-\mu}{\sigma},$$

V. imp

- *The probability density function for the normal distribution in standard form is given by*


$$f(Z) = \frac{1}{\sqrt{2\pi}}\, e^{-\frac{1}{2}(Z)^2}$$

- *Probability*

$$P(z_1 < z < z_2) = \int_{z_1}^{z_2} f(z)dz$$

©Gateway Classes : 7455 9612 81

Standard Normal Curve



Probability

$P(z_1 < z < z) =$ *Area under standard normal curve between* $z = z_1$ *and* $z = z_2$

Gateway Classes : 7455 9612 84

Q. 1 *If X is a normal variate with mean 80 and standard deviation 10, find probability $P(X \leq 100)$.*

Given

$\mu = 80$

$\sigma = 10$


$x = 100$

$$z = \frac{x - \mu}{\sigma}$$

$$z = \frac{100 - 80}{10}$$

$$\boxed{z = 2}$$

$$P(x \leq 100) = P(z \leq 2)$$



$$= P(-\infty < z < 0) + P(0 < z \leq 2)$$

$$= 0.5 + P(0 < z \leq 2)$$

$$= 0.5 + 0.4772$$

$$\boxed{P(x \leq 100) = 0.9772}$$

Gateway Classes 7455 961284

Q.2 *The life of army shoes is normally distributed with mean 8 months and standard deviation 2 months. If 5000 pairs are insured, how many pairs would be expected to need replacement after 12 months?* $\left[\text{Given that } P(z \geq 2) = 0.0228 \text{ and } z = \frac{x-\mu}{\sigma}\right]$. **(A.K.T.U. 2018)**

Given

$\mu = 8$

$\sigma = 2$

$N = 5000$

Let $x = 12$


$P(x \geq 12) = ?$

$$z = \frac{x-\mu}{\sigma}$$

$$= \frac{12-8}{2}$$

$$\boxed{z = 2}$$

$P(x \geq 12) = P(z \geq 2)$



$P(x \geq 12) = P(0 < z < \infty) - P(0 < z < 2)$

Gateway Classes : 7455 961284

$P(x \geq 12) = 0.5 - 0.4772$

$= 0.0228$

Number of pairs having life more then 12 months

$= 5000 \times 0.0228$

$= 114$

No. of pairs replaced after 12 months $= 5000 - 114$

$= 4886$

Gateway Classes 7455961284

Q. 3 *Assume mean height of soldiers to be 68.22 inches with a variance of 10.8 inches square. How many soldiers in a regiment of 1,000 would you expect to be over 6 feet tall, given that the area under the standard normal curve between $z = 0$ and $z = 0.35$ is 0.1368 and between $z = 0$ and $z = 1.15$ is 0.3746.* [AKTU (C.O.) 2021

Given

$\mu = 68.22$

$\sigma^2 = 10.8$

$\sigma = \sqrt{10.8}$

$N = 1000$

$x = 6$ feet

$x = 6 \times 12$


$x = 72$ inches

$z = \frac{x - \mu}{\sigma}$

$z = \frac{72 - 68.22}{\sqrt{10.8}}$

$z = 1.15$

$P(x > 72) = P(z > 1.15)$



GatewayClasses : 7455 9612 84

$$P(x > 72) = P(z > 1.15)$$

$$= P(0 < z < \infty) - P(0 < z < 1.15)$$

$$= 0.5 - 0.3746$$

$$= 0.1254$$

$$\text{No. of Soldiers} = 1000 \times 0.1254$$

$$= 125 \text{ (approx)}$$

Gateway Classes : 7455 9612 84

Q.4 *The daily wages of 1000 workers are distributed around a mean of Rs. 140 and with a standard deviation of Rs. 10. Estimate the number of workers whose daily wages*

(i) between 140 and 144 *(ii) less than 126* *(iii) more than 160.* **(AKTU. 2021)**

Given

$N = 1000$

$\mu = 140$

$\sigma = 10$

$$z = \frac{x - \mu}{\sigma}.$$

$$z = \frac{x - 140}{10} \quad \text{---} \quad (1)$$


(i) $x_1 = 140$, $x_2 = 144$

$$z_1 = \frac{140 - 140}{10} = 0$$

$z_1 = 0$

$$z_2 = \frac{144 - 140}{10} = \frac{4}{10}$$

$z_2 = 0.4$



$$P(140 < x < 144) = P(0 < z < 0.4)$$

$$= 0.1554$$

Gateway Classes : 7455 96 12 84

Expected number of workers

$= 0.1554 \times 1000$

$= 155.4$

$= 155$


(ii) $x = 126$

$$Z = \frac{x - 140}{10}$$

$$= \frac{126 - 140}{10}$$

$z = -1.4$

$P(x < 126) = P(z < -1.4)$



$P(x < 126) = P(z > 1.4)$

$= P(0 < z < \infty) - P(0 < z < 1.4)$

Gateway Classes: 7455 9612 84

$P(x<126) = 0.5 - 0.4192$

$= 0.0808$

Expected no. of workers

$= 1000 \times 0.0808$

$= 80.8$


$= 81$ (approx)

(iii) $x = 160$

$z = \frac{160 - 140}{10} = \frac{20}{10} = 2$

$z = 2$

$P(x > 160) = P(z > 2)$



$= P(0 < z < \infty) - P(0 < z < 2)$

$= 0.5 - 0.4772 = 0.0228$

Expected no. of workers

$= 0.0228 \times 1000 = 22.8$

$= 23$

Gateway Classes: 7455 9612 84

Topic : Normal Probability Distribution (Part – I)

Q. 1 Define:
(i) Binomial distribution
(ii) Poisson distribution
(iii) Normal distribution. (AKTU 2018)

Q. 2 If X is a normal variate with mean 30 and S. D. 5, find the probabilities that
(i) $26 \leq X \leq 40$
(ii) $X \geq 45$ and
(iii) $[X - 30 | > 5$

Ans. 2(i) 0.7653 (ii) 0.00135 (iii) 0.3174

Q. 3 If Z is a standard normal variable, find the following probabilities:

(i) $P(Z < 1.2)$
(ii) $P(Z > -1.2)$
(iii) $P(-1.2 < Z < 1.3)$.

Ans. 3(i) 0.8849 (ii) 0.8849 (iii) 0.7881

Gateway Classes : 7455 9612 84

ENGINEERING MATHEMATICS

UNIT-4 : *Statistical Techniques – II*

Lec-09

Today's Target


- $\boldsymbol{Normal\ Probability\ Distribution(Part-II)}$
- PYQs
- DPPs

Gateway Classes : 7455 961284

Standard Normal Curve

$-\infty < z < 0$

$0 < z < \infty$



$-\infty < z < \infty$

$$z = \frac{x - \mu}{\sigma},$$

Total Area under curve = 1

From Normal table

Probability

$P(z_1 < z < z_2)$ = **Area under standard normal Curve between** $z = z_1$ **and** $z = z_2$

Where

x = Normal random variable

Z = Standard normal random variable

μ = Mean

σ = Standard deviation

Gateway Classes : 7455 962 84

Q.1 *A sample of 100 dry battery cells tested to find the length of life produced the following results:* $\bar{x} = 12$ *hours,* $\sigma = 3$ *hours. Assuming the data to be normally distributed, what % of battery cells are expected to have life* (i) *more than 15 hours* (ii) *less than 6 hours* (iii) *between 10 and 14 hours?* (*A.K.T.U. 2018, 2021*)

Given

$N = 100$

$\mu = 12$

$\sigma = 3$

$$z = \frac{x - \mu}{\sigma}$$


$$z = \frac{x - 12}{3} \quad \text{—— (1)}$$

(i) $x = 15$

$$z = \frac{15 - 12}{3} = \frac{3}{3}$$

$z = 1$

$P(x \geq 15) = P(z > 1)$



$P(x > 15) = P(0 < z < \infty) - P(0 < z < 1)$

$= 0.5 - 0.3413$

$= 0.1587$

Gateway Classes : 7455 9612 84

$P(x>15) = 0.1587 \times 100$

$= 15.87\%$


(iii) $x = 6$

$$z = \frac{x-12}{3} = \frac{6-12}{3}$$

$z = -2$

$P(x<6) = P(z<-2)$

$= P(z>2)$



$P(x<6) = P(0<z<\infty) - P(0<z<2)$

$= 0.5 - 0.4772$

$= 0.0228$

$P(x<6) = 0.0228 \times 100 = 2.28\%$

Gateway Classes 7455 9612 84

$P(x<6) = 2.28\%$

(iii) $x_1 = 10$

$x_2 = 14$

$$z_1 = \frac{x_1 - 12}{3}$$


$$z_1 = \frac{10 - 12}{3}$$

$$z_1 = -0.67$$

$$z_2 = \frac{x_2 - 12}{3}$$

$$z_2 = \frac{14 - 12}{3}$$

$$z_2 = 0.67$$



$P(10 < x < 14) = P(-0.67 < z < 0.67)$

$= 2P(0 < z < 0.67)$

$= 2 \times 0.2485 = 0.497$

$P(10 < x < 14) = 49.7\%$

Gateway Classes : 7455961284

GW
GATEWAY CLASSES

Standard Normal C

z	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
0.0	.0000	.0040	.0080	.0120	.0160	.0199	.0239	.0279	.0319	.0359
0.1	.0398	.0438	.0478	.0517	.0557	.0596	.0636	.0675	.0714	.0754
0.2	.0793	.0832	.0871	.0910	.0948	.0987	.1026	.1064	.1103	.1141
0.3	.1179	.1217	.1255	.1293	.1331	.1368	.1406	.1443	.1480	.1517
0.4	.1554	.1591	.1628	.1664	.1700	.1736	.1772	.1808	.1844	.1879
0.5	.1915	.1950	.1985	.2019	.2054	.2088	.2123	.2157	.2190	.2224
0.6	.2257	.2291	.2324	.2357	.2389	.2422	.2454	.2485	.2517	.2549
0.7	.2580	.2611	.2642	.2673	.2704	.2734	.2764	.2794	.2823	.2852
0.8	.2881	.2910	.2939	.2967	.2995	.3023	.3051	.3078	.3106	.3133
0.9	.3159	.3186	.3212	.3238	.3264	.3289	.3315	.3340	.3365	.3389
1.0	.3413	.3438	.3461	.3485	.3508	.3531	.3554	.3577	.3599	.3621
1.1	.3643	.3665	.3686	.3708	.3729	.3749	.3770	.3790	.3810	.3830
1.2	.3849	.3869	.3888	.3907	.3925	.3944	.3962	.3980	.3997	.4015
1.3	.4032	.4049	.4066	.4082	.4099	.4115	.4131	.4147	.4162	.4177
1.4	.4192	.4207	.4222	.4236	.4251	.4255	.4279	.4292	.4306	.4319
1.5	.4332	.4345	.4357	.4370	.4382	.4394	.4406	.4418	.4429	.4441
1.6	.4452	.4463	.4474	.4484	.4495	.4505	.4515	.4525	.4535	.4545
1.7	.4554	.4564	.4573	.4582	.4591	.4599	.4608	.4616	.4625	.4633
1.8	.4641	.4649	.4656	.4664	.4671	.4678	.4686	.4693	.4699	.4706
1.9	.4713	.4719	.4726	.4732	.4738	.4744	.4750	.4756	.4761	.4767
2.0	.4772	.4778	.4783	.4788	.4793	.4798	.4803	.4808	.4812	.4817
2.1	.4821	.4826	.4830	.4834	.4838	.4842	.4846	.4850	.4854	.4857
2.2	.4861	.4864	.4868	.4871	.4875	.4878	.4881	.4884	.4887	.4890
2.3	.4893	.4896	.4898	.4901	.4904	.4906	.4909	.4911	.4913	.4916
2.4	.4918	.4920	.4922	.4925	.4927	.4929	.4931	.4932	.4934	.4936
2.5	.4938	.4940	.4941	.4943	.4945	.4946	.4948	.4949	.4951	.4952
2.6	.4953	.4955	.4956	.4957	.4959	.4930	.4961	.4962	.4963	.4964
2.7	.4965	.4966	.4967	.4968	.4969	.4970	.4971	.4972	.4973	.4974
2.8	.4974	.4975	.4976	.4977	.4977	.4978	.4979	.4979	.4980	.4981
2.9	.4981	.4982	.4982	.4983	.4984	.4984	.4985	.4985	.4986	.4986
3.0	.4987	.4987	.4987	.4988	.4988	.4989	.4999	.4989	.4990	.4990
3.1	.4990	.4991	.4991	.4991	.4992	.4992	.4992	.4992	.4993	.4993

Gateway Classes +4552982284

Q. 2 *In a sample of 1000 cases, the mean of a certain test is 14 and S.D. is 2.5.* **(A.K.T.U. 2020)**

Assuming the distribution to be normal, find

(i) how many students score between 12 and 15?

(ii) how many score above 18?

(iii) how many score below 8?

(iv) how many score 16?

Given

$\mu = 14$

$\sigma = 2.5$

$N = 1000$

$z = \frac{x - \mu}{\sigma}$

$z = \frac{x - 14}{2.5}$ ——— (1)

(i) $x_1 = 12$, $x_2 = 15$


$z_1 = \frac{x_1 - 14}{2.5} = \frac{12 - 14}{2.5}$

$z_1 = -0.8$

$z_2 = \frac{x_2 - 14}{2.5} = \frac{15 - 14}{2.5}$

$z_2 = 0.4$

$P(12 < x < 15) = P(-0.8 < z < 0.4)$



Gateway Classes : 7455 961 384

$P(12<n<15) = P(0<z<0.4) + P(0<z<0.8)$

$= 0.1554 + 0.2881$

$= 0.4435$

Number of students $= 1000 \times 0.4435$

$= 443.5$

$= 444$

(ii) $n = 18$


$z = \frac{n-14}{2.5} = \frac{18-14}{2.5} = \frac{4}{2.5}$

$z = 1.6$

$P(n > 18) = P(z > 1.6)$

$= P(0<z<\infty) - P(0<z<1.6)$

$= 0.5 - 0.4452$



Gateway Classes : 7455 961284

$P(x > 18) = 0.0548$

Number of students $= 1000 \times 0.0548$

$= 54.8$

$\boxed{= 55}$

(iii) $x = 8$


$$z = \frac{x - 14}{2.5}$$

$$z = \frac{8 - 14}{2.5}$$

$z = -2.4$

$P(x < \infty) = P(z < -2.4)$

$= P(z > 2.4)$



$P(x < 8) = P(0 < z < \infty) - P(0 < z < 2.4)$

$= 0.5 - 0.4918$

$= 0.0082$

Gateway Classes : 7455 961284

Number of students $= 0.0082 \times 1000$

$= 8.2$

$= 8$

(IV) $x = 16$

$x_1 = 15.5, \quad x_2 = 16.5$


$$z_1 = \frac{x_1 - 14}{2.5} = \frac{15.5 - 14}{2.5}$$

$$z_1 = 0.6$$

$$z_2 = \frac{x_2 - 14}{2.5} = \frac{16.5 - 14}{2.5}$$

$z_2 = 1$

$P(15.5 < x < 16.5) = P(0.6 < z < 1)$



$P(15.5 < x < 16.5) = P(0 < z < 1) - P(0 < z < 0.6)$

$= 0.3413 - 0.2257$

$= 0.1156$

Number of students $= 0.1156 \times 1000$


$= 116$

Gateway Classes : 7455 9612 84

Q.3 *In a normal distribution, 31% of the items are under 45 and 8% are over 64. Find the mean and standard deviation of the distribution. It is given that that if*

$f(t) = \frac{1}{\sqrt{2\pi}} \int_0^t e^{-\frac{1}{2}x^2} dx$ then f(0.5) = 0.19 and f(1.4) = 0.42. **(A.K.T.U. 2020)**

Let mean = μ, Standard deviation = σ



When $x_1 = -45$, $P(-45 < x < 0) = \frac{19}{100}$

$= 0.19$

$z_1 = -0.5$

When $x_2 = 64$, $P(0 < x < 64)$

$= \frac{42}{100} = 0.42$

$z_2 = 1.4$

Gateway Classes : 7455 961284

$$z = \frac{x - \mu}{\sigma}$$

$$z_1 = \frac{x_1 - \mu}{\sigma}$$

$$-0.5 = \frac{45 - \mu}{\sigma}$$

$$-0.5\sigma = 45 - \mu$$

$$-0.5\sigma + \mu = 45 \quad \text{—— (1)}$$

$$z_2 = \frac{x_2 - \mu}{\sigma}$$

$$1.4 = \frac{64 - \mu}{\sigma}$$

$$1.4\sigma = 64 - \mu$$

$$1.4\sigma + \mu = 64 \quad \text{—— (2)}$$

Subtract (1) and (2)

$$-0.5\sigma + \mu = 45$$

$$1.4\sigma + \mu = 64$$

$$-1.9\sigma = -19$$

$$\sigma = 10$$

$$\mu = 50$$

GW GATEWAY CLASSES

Gateway Classes : 7455 9612 84

GW GATEWAY CLASSES

Standard Normal C

z	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
0.0	.0000	.0040	.0080	.0120	.0160	.0199	.0239	.0279	.0319	.0359
0.1	.0398	.0438	.0478	.0517	.0557	.0596	.0636	.0675	.0714	.0754
0.2	.0793	.0832	.0871	.0910	.0948	.0987	.1026	.1064	.1103	.1141
0.3	.1179	.1217	.1255	.1293	.1331	.1368	.1406	.1443	.1480	.1517
0.4	.1554	.1591	.1628	.1664	.1700	.1736	.1772	.1808	.1844	.1879
0.5	.1915	.1950	.1985	.2019	.2054	.2088	.2123	.2157	.2190	.2224
0.6	.2257	.2291	.2324	.2357	.2389	.2422	.2454	.2485	.2517	.2549
0.7	.2580	.2611	.2642	.2673	.2704	.2734	.2764	.2794	.2823	.2852
0.8	.2881	.2910	.2939	.2967	.2995	.3023	.3051	.3078	.3106	.3133
0.9	.3159	.3186	.3212	.3238	.3264	.3289	.3315	.3340	.3365	.3389
1.0	.3413	.3438	.3461	.3485	.3508	.3531	.3554	.3577	.3599	.3621
1.1	.3643	.3665	.3686	.3708	.3729	.3749	.3770	.3790	.3810	.3830
1.2	.3849	.3869	.3888	.3907	.3925	.3944	.3962	.3980	.3997	.4015
1.3	.4032	.4049	.4066	.4082	.4099	.4115	.4131	.4147	.4162	.4177
1.4	.4192	.4207	.4222	.4236	.4251	.4255	.4279	.4292	.4306	.4319
1.5	.4332	.4345	.4357	.4370	.4382	.4394	.4406	.4418	.4429	.4441
1.6	.4452	.4463	.4474	.4484	.4495	.4505	.4515	.4525	.4535	.4545
1.7	.4554	.4564	.4573	.4582	.4591	.4599	.4608	.4616	.4625	.4633
1.8	.4641	.4649	.4656	.4664	.4671	.4678	.4686	.4693	.4699	.4706
1.9	.4713	.4719	.4726	.4732	.4738	.4744	.4750	.4756	.4761	.4767
2.0	.4772	.4778	.4783	.4788	.4793	.4798	.4803	.4808	.4812	.4817
2.1	.4821	.4826	.4830	.4834	.4838	.4842	.4846	.4850	.4854	.4857
2.2	.4861	.4864	.4868	.4871	.4875	.4878	.4881	.4884	.4887	.4890
2.3	.4893	.4896	.4898	.4901	.4904	.4906	.4909	.4911	.4913	.4916
2.4	.4918	.4920	.4922	.4925	.4927	.4929	.4931	.4932	.4934	.4936
2.5	.4938	.4940	.4941	.4943	.4945	.4946	.4948	.4949	.4951	.4952
2.6	.4953	.4955	.4956	.4957	.4959	.4930	.4961	.4962	.4963	.4964
2.7	.4965	.4966	.4967	.4968	.4969	.4970	.4971	.4972	.4973	.4974
2.8	.4974	.4975	.4976	.4977	.4977	.4978	.4979	.4979	.4980	.4981
2.9	.4981	.4982	.4982	.4983	.4984	.4984	.4985	.4985	.4986	.4986
3.0	.4987	.4987	.4987	.4988	.4988	.4989	.4999	.4989	.4990	.4990
3.1	.4990	.4991	.4991	.4991	.4992	.4992	.4992	.4992	.4993	.4993

Gateway Classes 7455967129

DPP- 09

Topic :*Normal Probability Distribution (Part – II)*

Q. 1 In a test on 2000 electric bulbs, it was found that the life of a particular make, was normally distributed with an average life of 2040 hours and S. D. of 60 hours, estimate the number of bulbs likely to burn for

(A. K. T. U. 2021)

(i) more than 2150 hours
(it) less than 1950 hours
(iii) more than 1920 hours but less than 2160 hours.

Ans. 1 (i) 67 (ii) 134 (iii) 1909

Q. 2 The marks X obtained in Mathematics by 1000 students are normally distributed with meas 78% and standard deviation 11% Determine:

Ans. 2(i) 138 (ii)63.92% (iii) Between 60 and 96

(i) how many students got marks above 90%?
(ii) What was the highest marks obtained by the lowest 10% of students?
(iii) Within what limits did the middle 90% of the students lie?

Q. 3 In a normal distribution exactly normal, 7% of the items are under 35 and 89% areunder 63.

What are the mean and standard deviation of the distribution?

Ans. 3 50.3 and 10.33

Gateway Classes : 7455961284

GATEWAY CLASSES

AKTU Full Courses (Paid)

Download Gateway Classes Application

From Google Play store

All Subjects

Link in Description

Gateway Classes : 7455961284

Thank You

GATEWAY CLASSES